

Einstein-aether theory, violation of Lorentz invariance, and metric-affine gravity

Christian Heinicke¹, Peter Baekler², Friedrich W. Hehl^{1,3}

¹*Institut für Theoretische Physik, Universität zu Köln, 50923 Köln, Germany*

²*Fachbereich Medien, Fachhochschule Düsseldorf, University of Applied Sciences,
Josef-Gockeln-Str. 9, 40474 Düsseldorf, Germany*

³*Department of Physics and Astronomy,
University of Missouri-Columbia, Columbia, MO 65211, USA*

(Dated: 2005-04-01, *aether19.tex*)

We show that the Einstein-aether theory of Jacobson and Mattingly (J&M) can be understood in the framework of the metric-affine (gauge theory of) gravity (MAG). We achieve this by relating the aether vector field of J&M to certain post-Riemannian *nonmetricity* pieces contained in an independent linear connection of spacetime. Then, for the aether, a corresponding geometrical curvature-square Lagrangian with a massive piece can be formulated straightforwardly. We find an exact spherically symmetric solution of our model.

PACS numbers: 03.50.Kk; 04.20.Jb; 04.50.+h

Keywords: Violation of Lorentz invariance, nonmetricity, gravity, connection, aether, metric-affine gravity

I. INTRODUCTION

In an attempt to violate Lorentz invariance locally, J&M [37] introduced an extra timelike 4-vector field u of unit length, see also [8, 12, 17, 18, 21, 26, 36, 46] and the earlier work of Gasperini [23, 24] and Kostelecky and Samuel [42]. This “aether field” is governed by a Lagrangian carrying a kinetic term $\sim (\nabla u)^2$ and a massive term $\sim u^2$. The aether field equation is of (massive) Yang-Mills type

$$\nabla H + \text{lower order term} \sim \ell u, \quad (1)$$

with $H \sim \nabla u$ as field momentum (excitation) and ℓ as some constant. The aether field u can be considered to be an analogue of a Yang-Mills potential B .

The purpose of our note is to point out that such an aether theory can be reconstructed quite naturally as a specific example within the framework of MAG [27, 30]. The reasons are as follows:

1. Since the aim of the aether theory of J&M is to violate local Lorentz invariance, one should abandon the (flat) Minkowski as well as the (curved) Riemannian spacetime, which are rigidly and locally Lorentz invariant, respectively, and look for a *post*-Riemannian structure of spacetime appropriate for such an approach. If one introduces in spacetime, besides the metric $g_{\alpha\beta}$, an independent linear connection 1-form $\Gamma_\alpha{}^\beta$, then it is known from literature, see [30], that the *nonmetricity*

$$Q_{\alpha\beta} := - \overset{\Gamma}{D} g_{\alpha\beta}, \quad (2)$$

where $\overset{\Gamma}{D}$ is the covariant exterior derivative with respect to the connection $\Gamma_\alpha{}^\beta$, *is a measure for the violation of local Lorentz invariance*. Note that the introduction of a torsion 2-form T^α is voluntary in this context.

2. In four dimensions the nonmetricity with its 40 components can be split irreducibly into four pieces (see [30]):

$$\begin{aligned} Q_{\alpha\beta} &= {}^{(1)}Q_{\alpha\beta} \oplus {}^{(2)}Q_{\alpha\beta} \oplus {}^{(3)}Q_{\alpha\beta} \oplus {}^{(4)}Q_{\alpha\beta}, \\ 40 &= 16 \oplus 16 \oplus 4 \oplus 4. \end{aligned} \quad (3)$$

In this article we want to concentrate on reconstructing the J&M theory in the framework of MAG. Therefore we pick the *two vector-like quantities* ${}^{(3)}Q_{\alpha\beta}$ and ${}^{(4)}Q_{\alpha\beta}$, with four independent components each, as relevant for our purpose. The explicit form of these two quantities will be studied below. A result will be that the geometrical interpretation of ${}^{(4)}Q_{\alpha\beta}$ identifies it as equivalent to the *Weyl covector* [84] of 1918, which is related to scale transformations (and thus extends the Lorentz to the conformal group), whereas ${}^{(3)}Q_{\alpha\beta}$ is related to *shear* transformations and, accordingly, is a generic field obstructing local Lorentz invariance. In other words, we identify ${}^{(3)}Q_{\alpha\beta}$ as analogue of the aether field u of J&M. Accordingly, by assuming an independent linear connection of spacetime, we arrive straightforwardly at a vector-like quantity of

shear type that dissolves local Lorentz invariance. It is part of the geometric structure of spacetime and represents as such some kind of genuine aether.

3. Identifying ${}^{(3)}Q_{\alpha\beta}$ as an analogue of the aether field u suggests a Yang-Mills type Lagrangian

$$V_{(3)Q} \sim \overset{\Gamma}{D} {}^{(3)}Q_{\alpha\beta} \wedge \star \overset{\Gamma}{D} {}^{(3)}Q^{\alpha\beta} + \ell {}^{(3)}Q_{\alpha\beta} \wedge \star {}^{(3)}Q^{\alpha\beta}, \quad (4)$$

where $\star$ denotes the Hodge star operator. The first term parallels the kinetic aether term of J&M, see [37], Eq.(2), the second one the massive term of J&M. If a gravitational Hilbert-Einstein Lagrangian is added to (4), we recover the basic structure of the J&M Lagrangian.

However, we stress an important difference. In our approach we proceed from variables of purely geometric origin: metric $g_{\alpha\beta}$ (defining angles and lengths), coframe ϑ^α (representing the local reference frame) and connection $\Gamma_\alpha{}^\beta$ (defining parallel displacement). In contrast, the aether field of J&M is an *external* quantity with no obvious relation to the geometry of spacetime, it is more like a cosmological fluid. Moreover, in spite of the presumed violation of Lorentz invariance, the J&M aether field u has a *fixed* (timelike) *magnitude*, which doesn't seem convincing to us.

In actual fact, the geometrical Lagrangian that we are going to study below will be a bit more complicated than (4). This conforms better to the MAG models that we developed earlier. In this context we will also demonstrate that the Lagrangian (4), in spite of the existence of the derivative terms in (4), fits very well into the first order version of MAG.

4. In the J&M model, the coupling of the aether field to matter seems to be an unsolved problem. In our approach, in the sense of Einstein's equivalence principle, we assume *minimal coupling*, that is, partial (or exterior) derivatives are substituted by covariant ones: $d \longrightarrow \overset{\Gamma}{D}$. Thereby a universal coupling of matter to our aether is guaranteed.

In Sec.II we give a sketch of metric-affine geometry, and in Sec.III we describe the main geometrical properties of the nonmetricity. In Sec.IV we turn to the the different curvature 2-forms, in particular to those that relate to post-Riemannian structures. Subsequently, in Sec.V, we display the general form of the field equations of MAG and compare them with those of J&M. In Sec.VI we formulate the general quadratic Lagrangian of MAG. We

tailor it such that it becomes somewhat analogous to the J&M Lagrangian. We compute the corresponding excitations (field momenta) explicitly by suitable partial differentiation of the Lagrangian. Accordingly, the field equations are now known explicitly.

In Sec.VII we present an exact spherically symmetric solution, the mass and the angular momentum of which are determined in Sec.VIII. The prolongation techniques that we took for finding exact solutions are explained shortly in Sec.IX. We discuss our results in Sec.X.

Our notation is as follows (see [30, 32]): We use the formalism of exterior differential forms. We denote the frame by e_α , with the anholonomic or frame indices $\alpha, \beta, \dots = 0, 1, 2, 3$. Decomposed with respect to a natural frame ∂_i , we have $e_\alpha = e^i_\alpha \partial_i$, where $i, j, \dots = 0, 1, 2, 3$ are holonomic or coordinate indices. The frame e_α is the vector basis of the tangent space at each point of the 4D spacetime manifold. The symbol $\lrcorner$ denotes the interior and $\wedge$ the exterior product. The coframe $\vartheta^\beta = e_j^\beta dx^j$ is dual to the frame, i.e., $e_\alpha \lrcorner \vartheta^\beta = \delta_\alpha^\beta$. If $\star$ denotes the Hodge star operator and if $\vartheta^{\alpha\beta} := \vartheta^\alpha \wedge \vartheta^\beta$, etc., then we can introduce the eta-basis $\eta := \star 1$, $\eta^\alpha := \star \vartheta^\alpha$, $\eta^{\alpha\beta} := \star \vartheta^{\alpha\beta}$, etc., see also [73]. Parentheses surrounding indices $(\alpha\beta) := (\alpha\beta + \beta\alpha)/2$ denote symmetrization and brackets $[\alpha\beta] := (\alpha\beta - \beta\alpha)/2$ antisymmetrization.

II. METRIC-AFFINE GEOMETRY OF SPACETIME

Spacetime is a 4-dimensional differentiable manifold that is equipped with a metric and a linear (also known as affine) connection (see [69]). The metric is of Lorentzian signature $(-+++)$ and is given by

$$g = g_{\alpha\beta} \vartheta^\alpha \otimes \vartheta^\beta. \quad (5)$$

In general, the coframe is left arbitrary, sometimes it is convenient to choose it orthonormal: $g = o_{\alpha\beta} \vartheta^\alpha \otimes \vartheta^\beta$, with $o_{\alpha\beta} = \text{diag}(-1, +1, +1, +1)$. The linear connection $\Gamma_\alpha{}^\beta$ governs parallel transfer and allows to define a covariant exterior derivative D (we drop now the Γ on top of D for convenience). We can decompose $\Gamma_\alpha{}^\beta$ with respect to a natural frame:

$$\Gamma_\alpha{}^\beta = \Gamma_{i\alpha}{}^\beta dx^i. \quad (6)$$

It has apparently 64 independent components. The notion of a non-trivial linear connection is decisive in going beyond the (flat) Minkowskian spacetime. To quote Einstein [16]: “... the essential achievement of general relativity, namely to overcome ‘rigid’ space (i.e., the inertial

frame), is *only indirectly* connected with the introduction of a Riemannian metric. The directly relevant conceptual element is the ‘displacement field’ (Γ_{ik}^l), which expresses the infinitesimal displacement of vectors. It is this which replaces the parallelism of spatially arbitrarily separated vectors fixed by the inertial frame (i.e., the equality of corresponding components) by an infinitesimal operation. This makes it possible to construct tensors by differentiation and hence to dispense with the introduction of ‘rigid’ space (the inertial frame). In the face of this, it seems to be of secondary importance in some sense that some particular Γ field can be deduced from a Riemannian metric ...”

The metric $g_{\alpha\beta}$ induces a Riemannian (or Levi-Civita) connection 1-form $\tilde{\Gamma}_\beta^\alpha$. In holonomic coordinates, it reads

$$\tilde{\Gamma}_i^j = \tilde{\Gamma}_{ki}^j dx^k, \quad \tilde{\Gamma}_{ki}^j := \frac{1}{2} g^{jl} (\partial_i g_{kl} + \partial_k g_{il} - \partial_l g_{ik}) . \quad (7)$$

The post-Riemannian part of the connection, the *distortion*

$$N_\alpha^\beta := \Gamma_\alpha^\beta - \tilde{\Gamma}_\alpha^\beta , \quad (8)$$

is a tensor-valued 1-form. Its 64 independent components describes the deviation from Riemannian geometry. In Einstein’s theory of gravity, general relativity (GR), spacetime is Riemannian, that is, the distortion $N_{\alpha\beta}$ vanishes.

TABLE I: Gauge fields and Bianchi identities in metric-affine gravity (MAG)

Potential	Field strength	Bianchi identity
metric $g_{\alpha\beta}$	nonmetricity $Q_{\alpha\beta} = -Dg_{\alpha\beta}$	zeroth $DQ_{\alpha\beta} = 2R_{(\alpha}{}^\mu g_{\beta)\mu}$
coframe ϑ^α	torsion $T^\alpha = D\vartheta^\alpha$	first $DT^\alpha = R_\mu{}^\alpha \wedge \vartheta^\mu$
connection Γ_α^β	curvature $R_\alpha{}^\beta = d\Gamma_\alpha^\beta - \Gamma_\alpha{}^\mu \wedge \Gamma_\mu{}^\beta$	second $DR_\alpha{}^\beta = 0$

There are two other measures for the deviation of a connection from its Levi-Civita part: The nonmetricity $Q_{\alpha\beta}$ of (2) and the torsion

$$T^\alpha := D\vartheta^\alpha , \quad (9)$$

see Table I. If one develops the covariant exterior derivative on the right-hand-side of (2) and substitutes the torsion of (9) suitably, then, after some algebra, one finds for the distortion

explicitly

$$N_{\alpha\beta} = -e_{[\alpha]}T_{\beta]} + \frac{1}{2}(e_{\alpha]}e_{\beta]}T_{\gamma})\vartheta^{\gamma} + (e_{[\alpha]}Q_{\beta]\gamma})\vartheta^{\gamma} + \frac{1}{2}Q_{\alpha\beta}. \quad (10)$$

Nonmetricity and torsion can be recovered from $N_{\alpha\beta}$ straightforwardly:

$$Q_{\alpha\beta} = 2N_{(\alpha\beta)}, \quad T^{\alpha} = N_{\beta}{}^{\alpha} \wedge \vartheta^{\beta}. \quad (11)$$

We call the (negative of the) torsion dependent piece of $N_{\alpha\beta}$ the *contortion* 1-form $K_{\alpha\beta}$. Like the torsion T^{α} , it has 24 independent components, and we have $T^{\alpha} = -K_{\beta}{}^{\alpha} \wedge \vartheta^{\beta}$.

The torsion has 3 irreducible pieces (see [30]), its totally antisymmetric piece (computer name **axitor**, 4 components),

$${}^{(3)}T^{\alpha} := -\frac{1}{3}e_{\alpha]}(\vartheta^{\beta} \wedge T_{\beta}) , \quad (12)$$

its trace (**trator**, 4 components)

$${}^{(2)}T^{\alpha} := \frac{1}{3}\vartheta^{\alpha} \wedge T \quad \text{with} \quad T := e_{\alpha]}T^{\alpha}, \quad (13)$$

and its tensor piece (**tentor**, 16 components)

$${}^{(1)}T^{\alpha} := T^{\alpha} - {}^{(2)}T^{\alpha} - {}^{(3)}T^{\alpha}. \quad (14)$$

For the Riemannian spacetime of GR, we have $T^{\alpha} = 0$ and $Q_{\alpha\beta} = 0$. If we relax the former constraint, $T^{\alpha} \neq 0$, we arrive at the Riemann-Cartan (RC) spacetime of the viable Einstein-Cartan theory of gravity, see [9, 27, 52, 75]. Since still $Q_{\alpha\beta} = 0$, such a RC-spacetime carries a metric-compatible connection and, accordingly, a length is invariant under parallel displacement. By the same token, a RC-spacetime is locally Lorentz invariant.

Haugan and Lämmerzahl [28], see also [44], argue that the presence of a torsion of spacetime violates local Lorentz invariance. Similarly, Kostelecky [41] and Bluhm and Kostelecky [11] attempt to violate Lorentz invariance already on the level of a RC-spacetime. According to our point of view, it is more natural to have a *nonvanishing* nonmetricity under such circumstances. Abandoning Lorentz invariance suggests the presence of nonmetricity, i.e., $Q_{\alpha\beta} \neq 0$. This is what we assume for the rest of our article.

A broad overview over the subject of violating Lorentz invariance has been given by Bluhm [10], see also the references given there. In experiment and in observation [45, 51, 55, 74] there is presently no evidence for Lorentz violations. Nevertheless, from a theoretical point of view (string theory, quantum gravity) a violation of Lorentz invariance is expected at some level, for certain models, see [2, 3], e.g..

III. SOME PROPERTIES OF THE NONMETRICITY

In the presence of nonmetricity $Q_{\alpha\beta} = -Dg_{\alpha\beta} \neq 0$, let us parallelly transport two vectors u and v from a point P along a curve with tangent vector c to a neighboring point Q . The scalar product of the two vectors $g(u, v)$ will change according to the the Lie derivative $\mathcal{L}_c g(u, v)$. With the gauge covariant Lie derivative of a form ψ (see [30, 57, 73]) $\mathbb{L}_c \psi := c \rfloor D\psi + D(c \rfloor \psi)$, we have

$$\mathcal{L}_c g(u, v) = \mathbb{L}_c g(u, v) = (\mathbb{L}_c g_{\alpha\beta}) u^\alpha v^\beta = c \rfloor (Dg_{\alpha\beta}) u^\alpha v^\beta, \quad (15)$$

since $\mathbb{L}_c u^\alpha = 0$ and $\mathbb{L}_c v^\alpha = 0$ because of parallel transfer. Thus,

$$\mathcal{L}_c g(u, v) = -(c \rfloor Q_{\alpha\beta}) u^\alpha v^\beta = -c \rfloor [\mathcal{Q}_{\alpha\beta} u^\alpha v^\beta + Q g(u, v)], \quad (16)$$

where

$$Q := \frac{1}{4} Q_\alpha{}^\alpha, \quad \mathcal{Q}_{\alpha\beta} := Q_{\alpha\beta} - Q g_{\alpha\beta} \quad (17)$$

are the trace (Weyl covector) and the traceless (shear) part of the nonmetricity. In the case of vanishing shear, the scalar product just changes by a factor and the light-cone is left intact under parallel transfer. Otherwise, with shear $\mathcal{Q}_{\alpha\beta}$, the angle between the vectors u and v does change. Hence, by admitting nonmetricity, we dissolve the local Lorentz invariance of a Riemann-Cartan spacetime. We depicted in Fig.1 how absolute parallelism, length, and angles are successively abandoned.

As we displayed in (3), the nonmetricity can be decomposed into four pieces. We have to recapitulate some of these features. In four dimensions, as symmetric tensor-valued 1-form, the nonmetricity has 40 independent components. Two vector-like pieces can be easily identified. Firstly, the Weyl covector $(17)_1$ can be extracted by taking the trace of $Q_{\alpha\beta}$. The remaining tracefree part of the nonmetricity $\mathcal{Q}_{\alpha\beta}$ in $(17)_2$ contains a second vector-like piece represented by the 1-form Λ :


$$\Lambda := (e^\beta \rfloor \mathcal{Q}_{\alpha\beta}) \wedge \vartheta^\alpha. \quad (18)$$

The 2-form¹

$$P_\alpha := \mathcal{Q}_{\alpha\beta} \wedge \vartheta^\beta - \frac{1}{3} \vartheta_\alpha \wedge \Lambda, \quad (19)$$

¹ For n dimensions, we have $P_\alpha := \mathcal{Q}_{\alpha\beta} \wedge \vartheta^\beta - \frac{1}{n-1} \vartheta_\alpha \wedge \Lambda$. In [30] we introduced a 2-form Ω_α instead that is related to P_α as follows: $P_\alpha = (-1)^{\text{Ind}(g)} \star \Omega_\alpha$.

FIG. 1: Two vectors at a point P span a triangle. If we parallelly transfer both vectors around a closed loop back to P , then in the course of the round trip the triangle gets linearly transformed.



with the properties

$$P^\alpha \wedge \vartheta_\alpha = 0, \quad e_\alpha \rfloor P^\alpha = 0, \quad (20)$$

turns out to be related to a further irreducible piece of $Q_{\alpha\beta}$. We have (see [30])

$${}^{(4)}Q_{\alpha\beta} := Q g_{\alpha\beta}, \quad (21)$$

$${}^{(3)}Q_{\alpha\beta} := \frac{4}{9} \left(\vartheta_{(\alpha} e_{\beta)} \rfloor - \frac{1}{4} g_{\alpha\beta} \right) \Lambda, \quad (22)$$

$${}^{(2)}Q_{\alpha\beta} := -\frac{2}{3} e_{(\alpha} \rfloor P_{\beta)}, \quad (23)$$

$${}^{(1)}Q_{\alpha\beta} := Q_{\alpha\beta} - {}^{(2)}Q_{\alpha\beta} - {}^{(3)}Q_{\alpha\beta} - {}^{(4)}Q_{\alpha\beta}. \quad (24)$$

As we want to relate the aether vector field u of J&M to the nonmetricity, it is obvious that ${}^{(4)}Q_{\alpha\beta} \sim Q$ and ${}^{(3)}Q_{\alpha\beta} \sim \Lambda$ are the objects of our main interest.

The nonmetricity $Q_{\alpha\beta} = Q_{i\alpha\beta} dx^i$, from a geometrical point of view, can be understood as a *strain* measure for the different directions specified by the 1-forms dx^i . In accordance with what we stated above, $(17)_2$ defines a shear measure $\mathcal{Q}_{\alpha\beta}$ since the dilation measure Q

is subtracted out. It is then immediately clear that ${}^{(4)}Q_{\alpha\beta} \sim Q$ is related to *dilations* and ${}^{(3)}Q_{\alpha\beta} \sim \Lambda$ to *shears*. Therefore, generically it is the 1-form Λ that is related to the aether vector u . However, we will keep also the Weyl 1-form Q since tentatively it seems to be related to the constraint $u^2 = 1$ of J&M.

Later, for the prolongation of previously known solutions, we need the expression $Q_{\alpha\beta} \wedge \vartheta^\beta$. It is useful to express it also in terms of Q , Λ , and P_α . If we substitute the irreducible decomposition (3) into $Q_{\alpha\beta} \wedge \vartheta^\beta$ and remember (21) to (24), we find

$$Q_{\alpha\beta} \wedge \vartheta^\beta = P_\alpha - \frac{1}{3} (\Lambda - 3Q) \wedge \vartheta_\alpha. \quad (25)$$

Note that ${}^{(1)}Q_{\alpha\beta} \wedge \vartheta^\beta = 0$, whereas the other irreducible pieces contribute.

As we saw in (4), we need “massive” terms in the nonmetricity for the construction of a J&M type Lagrangian. Since ${}^{*(4)}Q^{\alpha\beta} = g_{\alpha\beta} {}^*Q$, we find straightforwardly

$${}^{(4)}Q_{\alpha\beta} \wedge {}^{*(4)}Q^{\alpha\beta} = 4Q \wedge {}^*Q. \quad (26)$$

In the case of ${}^{(3)}Q_{\alpha\beta}$, things are a bit more complicated. However, if we use the formulas $\vartheta^\alpha \wedge {}^*\vartheta_\beta = \delta_\beta^\alpha \eta$ and $\eta = {}^*1$, then, after some algebra, we arrive at the same type of formula

$${}^{(3)}Q_{\alpha\beta} \wedge {}^{*(3)}Q^{\alpha\beta} = \frac{4}{9} \Lambda \wedge {}^*\Lambda. \quad (27)$$

We find the simplicity of (26) and (27) remarkable.

IV. CURVATURE

The standard definition of the curvature 2-form $R_\alpha{}^\beta$ is given in Table I. The Riemannian curvature (of GR) we denote by $\tilde{R}_\alpha{}^\beta$; of course, $\tilde{R}_{\alpha\beta} = -\tilde{R}_{\beta\alpha}$. The curvature $R_\alpha{}^\beta$ can be decomposed into its symmetric and antisymmetric part according to

$$R_{\alpha\beta} = W_{\alpha\beta} + Z_{\alpha\beta}, \quad \text{with} \quad W_{\alpha\beta} := R_{[\alpha\beta]}, \quad Z_{\alpha\beta} := R_{(\alpha\beta)}. \quad (28)$$

The “rotational” curvature $W_{\alpha\beta}$ characterizes a RC-space, whereas the “strain” curvature $Z_{\alpha\beta}$ only emerges if nonmetricity is admitted. Therefore, the investigation of the “Lorentz violating” curvature $Z_{\alpha\beta}$ is of central importance to our paper.

However, in order to link up our theory to GR and to the Einstein-Cartan theory with its RC-spacetime, we have to take a look at the rotational curvature $W_{\alpha\beta}$ as well. It can be

decomposed into six irreducible pieces:

$$\begin{aligned}
W_{\alpha\beta} &= {}^{(1)}W_{\alpha\beta} \oplus {}^{(2)}W_{\alpha\beta} \oplus {}^{(3)}W_{\alpha\beta} \oplus {}^{(4)}W_{\alpha\beta} \oplus {}^{(5)}W_{\alpha\beta} \oplus {}^{(6)}W_{\alpha\beta} \\
&= \text{weyl} \oplus \text{paircom} \oplus \text{pscalar} \oplus \text{ricsymf} \oplus \text{ricanti} \oplus \text{scalar}, \\
36 &= 10 \oplus 9 \oplus 1 \oplus 9 \oplus 6 \oplus 1.
\end{aligned} \tag{29}$$

The names are those that we use in our computer algebra programs [71] for the decomposition of $W_{\alpha\beta}$. In a RC-space, all six pieces in (29) are nonvanishing in general. If torsion $T^\alpha = 0$, then, as can be seen from the first Bianchi identity in Table I, **paircom**, **pscalar**, and **ricanti** vanish and we are left with the three pieces known from GR: The Weyl tensor **weyl**, the symmetric tracefree Ricci tensor **ricsymf**, and the curvature scalar **scalar**. Thereby the curvature reduces from the 36 independent components in a RC-space to 20 independent components in a Riemannian space, a result well-known from GR.

Let us come back to the strain curvature $Z_{\alpha\beta} = Z_{\beta\alpha}$. Obviously, it has one distinctive piece, namely its trace $Z := g^{\alpha\beta}Z_{\alpha\beta} = Z_\gamma{}^\gamma$. It should be noted that Z is related to a *premetric* quantity. In a space in which only a linear connection is specified, the curvature $R_\alpha{}^\beta$ can be contracted, $R_\gamma{}^\gamma$, even if a metric is not present. Thus $R_\gamma{}^\gamma$ and, as a consequence, also Z is rightfully called **dilcurv**, the part of the curvature related to dil(at)ations. This is an irreducible piece of $Z_{\alpha\beta}$ and we call it

$${}^{(4)}Z_{\alpha\beta} := \frac{1}{4} g_{\alpha\beta} Z. \tag{30}$$

Since on the level of the nonmetricity dilations are related to ${}^{(4)}Q_{\alpha\beta}$, we denoted the related curvature piece by the same number. In fact, the zeroth Bianchi identity in Table I, if contracted, yields $g^{\alpha\beta}DQ_{\alpha\beta} = 2Z_\gamma{}^\gamma = Z$. By partial integration, we find

$$Z = 2dQ \quad \text{or} \quad {}^{(4)}Z_{\alpha\beta} = \frac{1}{2} g_{\alpha\beta} dQ = \frac{1}{2} (D {}^{(4)}Q_{\alpha\beta} + Q_{\alpha\beta} \wedge Q). \tag{31}$$

Apparently, to the field strength Z , that is, to **dilcurv**, there belongs the potential Q , the Weyl covector.²

The tracefree part of the strain curvature

$$\mathcal{Z}_{\alpha\beta} := Z_{\alpha\beta} - \frac{1}{4} Z g_{\alpha\beta} \tag{32}$$

² Let us recall, it was **dilcurv** that Weyl [84] used in his unsuccessful unified field theory of 1918 to describe the electromagnetic field strength F — and Q was meant to be the electromagnetic potential A , see Goenner [25] and O’Raifeartaigh [61].

represents the shear curvature. In terms of $Z_{\alpha\beta}$, a further decomposition of $Z_{\alpha\beta}$ is possible. We have collected the results in Appendix A. We find a decomposition of $Z_{\alpha\beta}$ into five irreducible pieces:

$$\begin{aligned} Z_{\alpha\beta} &= {}^{(1)}Z_{\alpha\beta} \oplus {}^{(2)}Z_{\alpha\beta} \oplus {}^{(3)}Z_{\alpha\beta} \oplus {}^{(4)}Z_{\alpha\beta} \oplus {}^{(5)}Z_{\alpha\beta}, \\ 60 &= 30 \oplus 9 \oplus 6 \oplus 6 \oplus 9. \end{aligned} \quad (33)$$

Alerted by our results with respect to the nonmetricity, we expect that ${}^{(3)}Z_{\alpha\beta}$ with its six independent components may be related to ${}^{(3)}Q_{\alpha\beta}$ in a similar way as ${}^{(4)}Z_{\alpha\beta}$ is to ${}^{(4)}Q_{\alpha\beta}$, see (31). Symbolically, we expect

$$\text{Weyl 1-form } Q \sim {}^{(4)}Q \xrightarrow{d} {}^{(4)}Z \sim dQ, \quad (34)$$

$$\text{shear 1-form } \Lambda \sim {}^{(3)}Q \xrightarrow{d} {}^{(3)}Z \sim d\Lambda. \quad (35)$$

The rigorous form of relation (34) was already presented in (31). What about (35)? Well, life is a bit more complicated than (35) suggests. If we take the definition of ${}^{(3)}Z_{\alpha\beta}$ from (A4) and the zeroth Bianchi identity $DQ_{\alpha\beta} = 2Z_{\alpha\beta}$ from Table I, then, after some light algebra, we find

$${}^{(3)}Z_{\alpha\beta} = \frac{1}{3} (2\vartheta_{(\alpha} \wedge e_{\beta)}] - g_{\alpha\beta}) \hat{\Delta}, \quad (36)$$

with

$$\hat{\Delta} = \frac{1}{4} \vartheta^\alpha \wedge e_\beta] (DQ_{\alpha\beta} - g_{\alpha\beta} DQ) = \frac{1}{4} \vartheta^\alpha \wedge e_\beta] (D\varrho_{\alpha\beta} + Q \wedge \varrho_{\alpha\beta}). \quad (37)$$

Apparently, Eq.(37) turns out to be more complicated than we guessed in (35). We can only hope to find a formula of type (35), if we forbid certain pieces of the connection to occur.

In Appendix B we will show that under the conditions

$${}^{(2)}Q_{\alpha\beta} = 0, \quad {}^{(1)}T^\alpha = {}^{(3)}T^\alpha = 0, \quad (38)$$

we have, see (B15) and (36),

$$\hat{\Delta} \stackrel{(38)}{=} \frac{1}{6} d\Lambda \quad \text{or} \quad {}^{(3)}Z_{\alpha\beta} \stackrel{(38)}{=} \frac{1}{18} (2\vartheta_{(\alpha} \wedge e_{\beta)}] - g_{\alpha\beta}) d\Lambda. \quad (39)$$

Accordingly (35) turns out to be correct after all, provided the conditions (38) are met. Thus, modulo the conditions (38), our goal is reached of constructing a gauge Lagrangian à la (4) in terms of Λ and Q . For ${}^{(4)}Q_{\alpha\beta}$ we have, see (30) and (31),

$${}^{(4)}Z_{\alpha\beta} \wedge {}^\star {}^{(4)}Z^{\alpha\beta} = \frac{1}{4} Z \wedge {}^\star Z = dQ \wedge {}^\star dQ, \quad (40)$$

and for ${}^{(3)}Q_{\alpha\beta}$, using (C1) and (39)₁,

$${}^{(3)}Z_{\alpha\beta} \wedge {}^*{}^{(3)}Z^{\alpha\beta} = \frac{4}{3} \hat{\Delta} \wedge {}^*\hat{\Delta} \stackrel{(38)}{=} \frac{1}{27} d\Lambda \wedge {}^*d\Lambda. \quad (41)$$

V. CURRENTS AND FIELD EQUATIONS OF MAG, COMPARISON WITH J&M

Similar as J&M [37], we assume gravity and aether to exist, and then we couple them minimally to certain matter fields Ψ . However, in our case the gauge potentials $(g_{\alpha\beta}, \vartheta^\alpha, \Gamma_\alpha{}^\beta)$ represent ordinary gravity $(g_{\alpha\beta}, \vartheta^\alpha)$ as well as the image $(\Gamma_\alpha{}^\beta)$ of the J&M aether, and they are both part of the geometry of spacetime. The total first-order Lagrangian reads

$$L_{\text{tot}} = V(g_{\alpha\beta}, \vartheta^\alpha, Q_{\alpha\beta}, T^\alpha, R_\alpha{}^\beta) + L(g_{\alpha\beta}, \vartheta^\alpha, \Psi, D\Psi). \quad (42)$$

The independent variables of the action principle are $g_{\alpha\beta}$, ϑ^α , $\Gamma_\alpha{}^\beta$, and Ψ . The variation of the matter Lagrangian

$$\delta L = \frac{1}{2} \delta g_{\alpha\beta} \sigma^{\alpha\beta} + \delta \vartheta^\alpha \wedge \Sigma_\alpha + \delta \Gamma_\alpha{}^\beta \wedge \Delta^\alpha{}_\beta + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} \quad (43)$$

allows us identify the material currents coupled to the potentials as metric and canonical energy-momentum and as hypermomentum, respectively: $(\sigma_{\alpha\beta}, \Sigma_\alpha, \Delta^\alpha{}_\beta)$. The energy-momenta $\sigma_{\alpha\beta}$ and Σ_α are related to each other by a Belinfante-Rosenfeld type of relation. The hypermomentum splits in spin current $\oplus$ dilation current $\oplus$ shear current:

$$\Delta_{\alpha\beta} = \tau_{\alpha\beta} + \frac{1}{4} g_{\alpha\beta} \Delta^\gamma{}_\gamma + \widehat{\Delta}^\gamma{}_{\alpha\beta}, \quad \tau_{\alpha\beta} = -\tau_{\beta\alpha}, \quad \widehat{\Delta}^\gamma{}_{\alpha\beta} = \widehat{\Delta}^\gamma{}_{\beta\alpha}, \quad \widehat{\Delta}^\gamma{}_\gamma = 0. \quad (44)$$

The hypothetical shear current $\widehat{\Delta}^\gamma{}_{\alpha\beta}$ is discussed in [31, 56], see also the literature given there.

Our strategy is to leave open the explicit form of the gauge Lagrangian V for the time being and to introduce the excitations (or field momenta) of the gauge fields instead:

$$M^{\alpha\beta} = -2 \frac{\partial V}{\partial Q_{\alpha\beta}}, \quad H_\alpha = -\frac{\partial V}{\partial T^\alpha}, \quad H^\alpha{}_\beta = -\frac{\partial V}{\partial R_\alpha{}^\beta}. \quad (45)$$

These three *constitutive* laws, expressing the excitations in terms of the field strengths, characterize the physical properties of the spacetime continuum under consideration.

The field equations read [30]

$$DM^{\alpha\beta} - m^{\alpha\beta} = \sigma^{\alpha\beta} \quad (\text{zeroth}), \quad (46)$$

$$DH_\alpha - E_\alpha = \Sigma_\alpha \quad (\text{first}), \quad (47)$$

$$DH^\alpha{}_\beta - E^\alpha{}_\beta = \Delta^\alpha{}_\beta \quad (\text{second}), \quad (48)$$

$$\frac{\delta L}{\delta \Psi} = 0 \quad (\text{matter}). \quad (49)$$

If the second field equation (48) is fulfilled, then either the zeroth field equation (46) or the first one (47) is redundant due to some Noether identities. Hence we need only to consider (46),(48),(49) or (47),(48),(49).

On the right-hand-side of each of the gauge field equations (46) to (48) we have a material current, on the left-hand-side first the Yang-Mills type term “derivative of excitation” minus, as second term, a gauge current that, together with the material current, features as source of the corresponding gauge field. The gauge currents turn out to be [30] the metrical energy-momentum of the gauge fields

$$m^{\alpha\beta} := 2 \frac{\partial V}{\partial g_{\alpha\beta}} = \vartheta^{(\alpha} \wedge E^{\beta)} + Q^{(\alpha}{}_\gamma \wedge M^{\beta)\gamma} - T^{(\alpha} \wedge H^{\beta)} - R_\gamma^{(\alpha} \wedge H^{|\gamma|\beta)} + R^{(\alpha|\gamma|} \wedge H^{\beta)}{}_\gamma, \quad (50)$$

the canonical energy-momentum of the gauge fields³

$$E_\alpha := \frac{\partial V}{\partial \vartheta^\alpha} = e_\alpha \rfloor V + (e_\alpha \rfloor T^\beta) \wedge H_\beta + (e_\alpha \rfloor R_\beta{}^\gamma) \wedge H^\beta{}_\gamma + \frac{1}{2}(e_\alpha \rfloor Q_{\beta\gamma}) M^{\beta\gamma}, \quad (51)$$

and the hypermomentum of the gauge fields

$$E^\alpha{}_\beta := \frac{\partial V}{\partial \Gamma_\alpha{}^\beta} = -\vartheta^\alpha \wedge H_\beta - g_{\beta\gamma} M^{\alpha\gamma}, \quad (52)$$

respectively.

Like J&M, we will concentrate on the sourcefree region, that is, we assume that the material currents vanish. We discussed them here in order to find the physical interpretations of the gauge currents $m^{\alpha\beta}$, E_α , and $E^\alpha{}_\beta$. Since J&M consider a symmetric energy-momentum in their theory, we consider the sourcefree zeroth and second field equations:

$$DM^{\alpha\beta} = m^{\alpha\beta}, \quad (53)$$

$$DH^\alpha{}_\beta = E^\alpha{}_\beta. \quad (54)$$

³ For the relations between different energy-momentum currents in gravitational theory one should also compare Itin [35].

These are the two field equations that underlie our model of the J&M theory. The rest of the paper will be devoted to making them explicit and for finding exact solutions of them.

The field equation (53) is of an Einsteinian type. If the gauge Lagrangian V depends on a Hilbert-Einstein term R_{sc} , inter alia (curvature scalar), then, within $m^{\alpha\beta}$, the Einstein 3-form emerges. The left hand side of (53) should depend on the shear 1-form Λ . This can be achieved by putting $M^{\alpha\beta} = -2\partial V/\partial Q_{\alpha\beta}$ proportional to ${}^{(3)}Q^{\alpha\beta}$, see (22). As a consequence, the Lagrangian V carries a quadratic ${}^{(3)}Q$ piece and we expect $V \sim R_{\text{sc}} + {}^{(3)}Q^2$. Under these circumstances, Eq.(53) is the analogue of [37], Eq.(5). Both equations have 10 independent components.⁴

Our second field equation (54), with the gauge hypermomentum as source, has 64 independent components. For this reason, as we argued above, we have to kill all components apart from those 4 components related to the shear 1-form $\Lambda \sim {}^{(3)}Q_{\alpha\beta}$. At a first glance, this seems to be an impossible task. A little reflection shows that the situation is not hopeless at all. Substitute (52) into (54):

$$DH^\alpha{}_\beta = -\vartheta^\alpha \wedge H_\beta - M^\alpha{}_\beta. \quad (55)$$

According to the the last paragraph, we have $M^{\alpha\beta} \sim {}^{(3)}Q^{\alpha\beta}$, that is, $M^\alpha{}_\beta$ on the right hand side of (55) depends on 4 independent components. The term with $H_\alpha = -\partial V/\partial T^\alpha$ can be chosen to vanish by forbidding explicit torsion dependent terms in the Lagrangian.

Left over for discussion is $H^\alpha{}_\beta = -\partial V/\partial R_\alpha{}^\beta$ on the left hand side of (55). Clearly we want this term to depend in an essential way on the shear 1-form Λ . A look at (41) convinces us to take $H^\alpha{}_\beta \sim {}^{(3)}Z^\alpha{}_\beta \sim d\Lambda$. Then the left hand side of (55) becomes a wave type expression $\sim \square\Lambda$ with four essential components. Hence our equation reduces to just four components, like the corresponding J&M equation. Thus, our second field equation (54) is undoubtedly the analogue of [37], Eq.(4). At the same time it is also clear that we could introduce a richer aether structure than the one J&M studied by means of their vector field u .

⁴ In our approach we don't find the analogue of the $\nabla(Ju)$ terms in the first line of the aether stress tensor of J&M [37], Eq.(9). The reason is clear. In the J&M aether these terms arise, see [17], p.2, "from varying the metric dependence of the connection." However, in our case the connection is an independent variable. Incidentally, an aether stress tensor that depends, as in the J&M theory, on second derivatives of the field variable is not particularly plausible anyway.

If we collect our heuristic arguments for constructing the MAG analogue of the J&M aether, then we arrive at

$$V_{\text{J\&M}} \sim \frac{1}{\kappa} (R_{\text{sc}} + {}^{(3)}Q^2) + \frac{1}{\rho} {}^{(3)}Z^2, \quad (56)$$

with κ as gravitational constant and ρ as a dimensionless coupling constant. This first toy Lagrangian should be compared with [37], Eq.(1). Our massive term ${}^{(3)}Q^2$ resemble the constraint piece in the J&M Lagrangian, whereas

$${}^{(3)}Z^2 \sim d\Lambda \wedge {}^*d\Lambda \sim g^{\gamma[\alpha} g^{\beta]\delta} (\partial_\alpha \Lambda_\beta) (\partial_\gamma \Lambda_\delta) \eta \quad (57)$$

is the analogue of $K^{ab}_{mn} \nabla_a u^m \nabla_b u^n$. However, J&M have, in K^{ab}_{mn} , four open constants c_1, c_2, c_3, c_4 . In the specific model investigated below, we concentrate on the simple Maxwell-type kinetic term (57). In doing so, we seem closer to Kostelecky and Samuel [42] than to J&M. However, the general MAG Lagrangian, see Appendix D, Eq.(D1), encompasses kinetic terms of all 11 irreducible pieces of the curvature and thereby generalizes the 4 parameters of J&M considerably.

Let us look back to our first ansatz for an aether Lagrangian in Eq.(4). There we had derivative terms of the nonmetricity. However, such terms are not allowed in first order MAG, see the gauge Lagrangian V in (42). Only an algebraic dependency of the field strengths $Q_{\alpha\beta}, T^\alpha, R_\alpha{}^\beta$, mostly quadratic for dimensional reasons, is allowed. Nevertheless, by the zeroth Bianchi identity, see Table I, the derivatives can be removed and transformed to terms algebraic in the curvature; this is at least possible for ${}^{(3)}Q_{\alpha\beta}$ and ${}^{(4)}Q_{\alpha\beta}$. Thus also the Lagrangian (4) falls into the category of allowed Lagrangians within MAG. At the same time we see again how closely nonmetricity and shear curvature are interwoven.

We would like to stress that the formalism of MAG that we developed in this paper up to now is exact and free of any hand waving arguments. It is a first order Lagrange-Noether gauge formalism of non-Abelian nature and our gauge field equations (53) and (54) are coupled nonlinear partial differential equations of second order in the gauge potentials $g_{\alpha\beta}, \Gamma_\alpha{}^\beta$. The only hand waving is involved in the *explicit* choice of the gauge Lagrangian $V(g_{\alpha\beta}, \vartheta^\alpha, Q_{\alpha\beta}, T^\alpha, R_\alpha{}^\beta)$ in (42). Since we want to develop a model that is an image of the J&M theory, we have to hand pick a suitable V . In (56) we made a first attempt.

VI. A LAGRANGIAN FOR GRAVITY AND AETHER

In gauge theories the Lagrangian is assumed to be quadratic in the field strengths, in our case in $R_\alpha{}^\beta$, T^α , and $Q_{\alpha\beta}$. The most general parity conserving Lagrangian of such a type has been displayed in Appendix D. For modeling the J&M aether theory, we don't need this very complicated expression in its full generality. Nevertheless, let us look at its basic structure:

$$V_{\text{MAG}} \sim \frac{1}{\kappa}(R_{\text{sc}} + \lambda_0 + T^2 + TQ + Q^2) + \frac{1}{\rho}(W^2 + Z^2). \quad (58)$$

All indices are suppressed. The expression in the first parentheses describes $(g_{\alpha\beta}, \vartheta^\alpha)$ -gravity of the Newton-Einstein type, including a cosmological term with λ_0 . This “weak” gravity is governed by the conventional gravitational constant κ . If only these terms are present, the propagation of $\Gamma_\alpha{}^\beta$ is inhibited. Then, in addition to conventional gravity, only new *contact* interactions emerge that are glued to matter. The *viable* Einstein-Cartan theory with its spin-spin contact interaction is an example.

If one desires to make the connection $\Gamma_\alpha{}^\beta$ propagating, see [43, 70] for vanishing and [4, 14, 20, 80, 81, 82, 83] for nonvanishing nonmetricity, then one has to allow for curvature-square pieces W^2 and Z^2 , as shown in (58). This “strong gravity”, the potential of which is $\Gamma_\alpha{}^\beta$, is governed by a new dimensionless coupling constant ρ . Our hypothesis is that such a universal strong gravitational interaction is present in nature.

The J&M aether, in our interpretation, allows at least the ${}^{(3)}Q_{\alpha\beta}$ piece of the connection to propagate, see the ${}^{(3)}Z^2$ piece in the ansatz (56). We would like to stick to (56) as closely as possible. Since we search for an exact spherically symmetric solution of our model to be defined, we need to be flexible in the exact choice of the Lagrangian. After some computer algebra *experiments*, we came up with the following toy Lagrangian that we are going to investigate in the context of the J&M aether theory:

$$V = \frac{1}{2\kappa} [-a_0 (R^{\alpha\beta} \wedge \eta_{\alpha\beta} + 2\lambda_0 \eta) + Q_{\alpha\beta} \wedge {}^\star (b_1 {}^{(1)}Q^{\alpha\beta} + b_3 {}^{(3)}Q^{\alpha\beta})] - \frac{z_3}{2\rho} R^{\alpha\beta} \wedge {}^\star {}^{(3)}Z_{\alpha\beta}. \quad (59)$$

We have the following constants of order unity: a_0, b_1, b_3, z_3 . As compared to (56), we have no torsion piece. However, we added in a ${}^{(1)}Q^{\alpha\beta}$ piece. The most general quadratic Lagrangian (D1) is appreciably more complicated than (59). Nevertheless, without our computer algebra programs (see [71], [33], [32]) we would not have been able to handle the messy expressions.

Once the Lagrangian is specified, it is simple to calculate the gauge excitations by *partial* differentiation of (59) with respect to $Q_{\alpha\beta}$, T^α , and $R_\alpha{}^\beta$:

$$M^{\alpha\beta} = -\frac{2}{\kappa} \star (b_1 {}^{(1)}Q^{\alpha\beta} + b_3 {}^{(3)}Q^{\alpha\beta}) , \quad (60)$$

$$H_\alpha = 0 , \quad (61)$$

$$H^\alpha{}_\beta = \frac{a_0}{2\kappa} \eta^\alpha{}_\beta + \frac{z_3}{\rho} \star {}^{(3)}Z^\alpha{}_\beta . \quad (62)$$

These excitations have to be substituted into the field equations (53),(54) and into the gauge currents (50),(51),(52), respectively.

VII. A SIMPLE SPHERICALLY SYMMETRIC SOLUTION OF MAG

We look for exact spherically symmetric solutions of the field equations belonging to the Lagrangian (59). For this purpose, the coframe ϑ^α is assumed to be of Schwarzschild-deSitter (or Kottler) form,

$$\vartheta^0 = e^{\mu(r)} dt , \quad \vartheta^1 = e^{-\mu(r)} dr , \quad \vartheta^2 = r d\theta , \quad \vartheta^3 = r \sin\theta d\phi , \quad (63)$$

with the function

$$e^{2\mu(r)} = 1 - 2\frac{m}{r} - \frac{\lambda_0}{3} r^2 . \quad (64)$$

We use Schwarzschild coordinates $x^i = (t, r, \theta, \phi)$. Since the coframe is assumed to be *orthonormal*, the metric reads

$$g = -\vartheta^0 \otimes \vartheta^0 + \vartheta^1 \otimes \vartheta^1 + \vartheta^2 \otimes \vartheta^2 + \vartheta^3 \otimes \vartheta^3 . \quad (65)$$

The nonmetricity $Q^{\alpha\beta}$ is given by

$$\begin{aligned} Q^{\alpha\beta} &= \frac{\ell_0 e^{-\mu(r)}}{2r^2} \begin{pmatrix} \vartheta^1 & 0 & 0 & 0 \\ 0 & 0 & \vartheta^2 & \vartheta^3 \\ 0 & \vartheta^2 & 0 & 0 \\ 0 & \vartheta^3 & 0 & 0 \end{pmatrix} + \frac{\ell_1 e^{-\mu(r)}}{2r^2} \begin{pmatrix} 3\vartheta^0 & 0 & -\vartheta^2 & -\vartheta^3 \\ 0 & -(\vartheta^0 - 3\vartheta^1) & 0 & 0 \\ -\vartheta^2 & 0 & 0 & 0 \\ -\vartheta^3 & 0 & 0 & 0 \end{pmatrix} \\ &= \frac{e^{-\mu(r)}}{2r^2} \begin{pmatrix} 3\ell_1 \vartheta^0 + \ell_0 \vartheta^1 & 0 & -\ell_1 \vartheta^2 & -\ell_1 \vartheta^3 \\ 0 & -\ell_1 (\vartheta^0 - 3\vartheta^1) & \ell_0 \vartheta^2 & \ell_0 \vartheta^3 \\ -\ell_1 \vartheta^2 & \ell_0 \vartheta^2 & 0 & 0 \\ -\ell_1 \vartheta^3 & \ell_0 \vartheta^3 & 0 & 0 \end{pmatrix} . \end{aligned} \quad (66)$$

The integration constants ℓ_0 and ℓ_1 can be interpreted as a measure for the violation of Lorentz invariance. According to (66), we have ${}^{(2)}Q^{\alpha\beta} = 0$ or $P_\alpha = 0$. All other irreducible pieces of $Q_{\alpha\beta}$ are nonvanishing: ${}^{(1)}Q^{\alpha\beta} \neq 0$, ${}^{(3)}Q^{\alpha\beta} \neq 0$, ${}^{(4)}Q^{\alpha\beta} \neq 0$. In particular, we find for the *shear* and the *Weyl* 1-forms

$$\Lambda = \frac{9e^{-\mu(r)}}{8r^2} (\ell_0 + \ell_1) \vartheta^1, \quad (67)$$

$$Q = \frac{e^{-\mu(r)}}{8r^2} [-4\ell_1 \vartheta^0 + (3\ell_1 - \ell_0) \vartheta^1]. \quad (68)$$

The torsion 2-form turns out to be

$$T^\alpha = \ell_0 \frac{e^{-\mu(r)}}{4r^2} \begin{pmatrix} \vartheta^{01} \\ 0 \\ -\vartheta^{12} \\ -\vartheta^{13} \end{pmatrix} - \ell_1 \frac{e^{-\mu(r)}}{4r^2} \begin{pmatrix} 0 \\ \vartheta^{01} \\ \vartheta^{02} \\ \vartheta^{03} \end{pmatrix} = \frac{e^{-\mu(r)}}{4r^2} \begin{pmatrix} \ell_0 \vartheta^{01} \\ -\ell_1 \vartheta^{01} \\ -\ell_1 \vartheta^{02} - \ell_0 \vartheta^{12} \\ -\ell_1 \vartheta^{03} - \ell_0 \vartheta^{13} \end{pmatrix}. \quad (69)$$

As a consequence, ${}^{(1)}T^\alpha = {}^{(3)}T^\alpha = 0$ and only ${}^{(2)}T^\alpha \neq 0$. By contraction of (69) with e_α we find (recall $T = e_\alpha T^\alpha$)

$$T = \frac{3e^{-\mu(r)}}{4r^2} (\ell_1 \vartheta^0 + \ell_0 \vartheta^1). \quad (70)$$

The requirement that the torsion (69) and the nonmetricity (66) together with the orthonormal coframe field (63) be a solution of the field equations of the Lagrangian (59) implies some constraints on the coupling constants. If we take care of these constraints in (59), we find the truncated Lagrangian

$$V = \frac{1}{2\kappa} \left[-R^{\alpha\beta} \wedge \eta_{\alpha\beta} - 2\lambda_0 \eta + Q_{\alpha\beta} \wedge \star \left(\frac{1}{4} {}^{(1)}Q^{\alpha\beta} - \frac{1}{2} {}^{(3)}Q^{\alpha\beta} \right) \right] - \frac{z_3}{2\rho} {}^{(3)}Z^{\alpha\beta} \wedge \star {}^{(3)}Z_{\alpha\beta}. \quad (71)$$

In other words, our exact solution (63)-(66),(69) solves the field equations of the Lagrangian (71). Note that z_3 is left arbitrary.

The explicit expressions for the strain and the rotational curvature can be found in⁵ Appendix F. The Weyl and the scalar parts of the rotational curvature $W_{\alpha\beta}$, see (F1) and (F6), are composed of a Riemannian and a post-Riemannian piece. The other parts of $W_{\alpha\beta}$ as well as of $Z_{\alpha\beta}$ are purely post-Riemannian. We were surprised that ${}^{(3)}Z^{\alpha\beta} = 0$; particularly simple is $\text{dilcurv} \sim \ell_1/(2r^3)$.

⁵ According to the classification scheme of Baekler et al. [7], this solution is a special subcase of class Va.

The set of the three one-forms $\{Q, T, \Lambda\}$ are related by

$$3Q + 2T - \Lambda = 0, \quad (72)$$

as can be checked easily. In this way, the torsion 1-form is closely related to the shear and the Weyl 1-forms. This is a relation which follows from Baekler's general prolongation ansatz [5] to solve the field equations of MAG.

Our solution looks like a superposition of two elementary solutions. For $\ell_1 = 0$ we find one elementary solution, a second one for $\ell_0 = 0$, both for the same set of coupling constants. Also this fact can be understood from the point of view of the prolongation technique.

VIII. KILLING VECTORS AND QUASILOCAL CHARGES

Let us determine the mass and the angular momentum of our exact solution. In a Riemannian space we call $\xi = \xi^\alpha e_\alpha$ a Killing vector if the latter is the generator of a symmetry transformation of the metric, i.e.,

$$\mathcal{L}_\xi g = 0. \quad (73)$$

In metric-affine space, coframe and connection are independent. Hence, Eq.(73) has to be supplemented by a corresponding requirement for the connection [30, p.83],

$$\mathcal{L}_\xi \Gamma_\alpha^\beta = 0. \quad (74)$$

These two relations can be recast into a more convenient form,

$$e_{(\alpha} \rfloor \tilde{D} \xi_{\beta)} = 0, \quad (75)$$

$$D \left(e_\alpha \rfloor \widehat{D} \xi^\beta \right) + \xi \rfloor R_\alpha^\beta = 0, \quad (76)$$

where $\tilde{D}$ refers to the Riemannian part of the connection (Levi-Civita connection) and $\widehat{D}$ to the transposed connection: $\widehat{D} := d + \widehat{\Gamma}_\alpha^\beta := d + \Gamma_\alpha^\beta + e_\alpha \rfloor T^\beta$.

For our solution the Killing vectors are the same as in case of the Schwarzschild-de Sitter metric in Riemannian spacetime,

$$\stackrel{(0)}{\xi} = \partial_t, \quad (77)$$

$$\stackrel{(1)}{\xi} = \sin \phi \partial_\theta + \cot \theta \cos \phi \partial_\phi, \quad (78)$$

$$\stackrel{(2)}{\xi} = -\cos \phi \partial_\theta + \cot \theta \sin \phi \partial_\phi, \quad (79)$$

$$\stackrel{(3)}{\xi} = \partial_\phi. \quad (80)$$

Subsequently we can compute the quasilocal charges by using formulas of Nester, Chen, Tung, and Wu [13, 34, 85, 86, 87], for related work see [6, 53], e.g. The barred quantities refer to a background solution, the symbol Δ denotes the difference between a solution and the background, $\Delta\alpha = \alpha - \alpha$.

$$\begin{aligned} \mathfrak{B}(N) := & - \left\{ \begin{array}{l} \frac{1}{2} \Delta g_{\alpha\beta} (N \rfloor \bar{M}^{\alpha\beta}) \\ \frac{1}{2} \Delta g_{\alpha\beta} (N \rfloor M^{\alpha\beta}) \end{array} \right\} - \left\{ \begin{array}{l} (N \rfloor \vartheta^\alpha) \Delta H_\alpha + \Delta \vartheta^\alpha \wedge (N \rfloor \bar{H}_\alpha) \\ (N \rfloor \bar{\vartheta}^\alpha) \Delta H_\alpha + \Delta \vartheta^\alpha \wedge (N \rfloor H_\alpha) \end{array} \right\} \\ & - \left\{ \begin{array}{l} (\widehat{D}_\alpha N^\beta) \Delta H^\alpha{}_\beta + \Delta \Gamma_\alpha{}^\beta \wedge (N \rfloor \bar{H}^\alpha{}_\beta) \\ \overline{(\widehat{D}_\alpha N^\beta)} \Delta H^\alpha{}_\beta + \Delta \Gamma_\alpha{}^\beta \wedge (N \rfloor H^\alpha{}_\beta) \end{array} \right\}. \end{aligned} \quad (81)$$

The upper (lower) line in the braces is chosen if the field strengths (momenta) are prescribed on the boundary. By taking $N = \partial_t$ and integrating $\mathfrak{B}$ over a 2-sphere and performing the limit $r \rightarrow \infty$, we get the total energy. As background solution we assume our solution with $m = 0$ and $\ell_0 = \ell_1 = 0$. Similarly, by taking $N = \partial_\phi$ we obtain the total angular momentum,

$$E_\infty = \lim_{r \rightarrow \infty} \int_{S^2} \mathfrak{B}(\partial_t) = -\frac{8\pi m}{\kappa}, \quad (82)$$

$$L_\infty = \lim_{r \rightarrow \infty} \int_{S^2} \mathfrak{B}(\partial_\phi) = 0. \quad (83)$$

IX. REMARKS ON THE PROLONGATION TECHNIQUE

Previously numerous exact solutions of MAG have already been found. Let us mention, as examples, the papers [29, 34, 39, 54, 59, 65, 68, 76, 77, 79, 82, 83, 85, 86] and references given there. Even possible links to observation were discussed in [1, 63, 64, 66, 67, 72]. The solution in Sec.VII was found by using prolongation methods. Such a method for MAG was proposed by Baekler et al. [5, 7]. In the sequel, we will explain how we applied the prolongation method to our case in question.

For this purpose, we start, in the framework of the Poincaré gauge theory (see [9, 27]), from a known exact solution with Schwarzschild metric and $1/r^2$ -torsion in a RC-spacetime, i.e., the nonmetricity vanishes. Then, for *generating* nonmetricity, we make the ansatz (Weyl 1-form $Q = Q_\alpha{}^\alpha/4$)

$$\begin{aligned} T^\alpha &= \xi_0 Q^\alpha{}_\beta \wedge \vartheta^\beta + \xi_1 Q \wedge \vartheta^\alpha + {}^{(3)}T^\alpha \\ &= \xi_0 Q^\alpha{}_\beta \wedge \vartheta^\beta + (\xi_0 + \xi_1) Q \wedge \vartheta^\alpha + {}^{(3)}T^\alpha, \\ 24 &= \quad 16 \quad + \quad 4 \quad + \quad 4, \end{aligned} \quad (84)$$

with arbitrary constants ξ_0 and ξ_1 to be determined by the field equations. The 24 components of T^α are related to the 40 components of $Q_{\alpha\beta}$. Because of the property ${}^{(1)}Q^\alpha_\beta \wedge \vartheta^\beta = 0$, the irreducible part ${}^{(1)}Q^\alpha_\beta$ (16 independent components) does not contribute to the torsion. Furthermore $(Q^\alpha_\beta \wedge \vartheta^\beta) \wedge \vartheta_\alpha = 0$ (4 independent components). Hence, our ansatz relates the 24 components of the torsion to $16 + 4 + 4 = 40 - 16$ independent components of $Q^\alpha_\beta \wedge \vartheta^\beta$, $Q \wedge \vartheta^\alpha$, and ${}^{(3)}T^\alpha$. In the end, the second order partial differential equations (53),(54) become nonlinear algebraic equations. We solve them by requiring certain *constraints* on the coupling constants. In this way we find exact solutions with $1/r^\nu$ -behavior of $Q_{\alpha\beta}$, here $\nu = 1, 2, 3$.

If we substitute the decomposition formula (25) into (84), we find

$$T^\alpha = \xi_0 P^\alpha - \frac{1}{3} [\xi_0 \Lambda - 3(\xi_0 + \xi_1) Q] \wedge \vartheta^\alpha + {}^{(3)}T^\alpha. \quad (85)$$

Contraction yields

$$\xi_0 \Lambda - 3(\xi_0 + \xi_1) Q - T = 0. \quad (86)$$

Here it is useful to take recourse to the first Bianchi identity. Provided that the conditions

$$T^\alpha = {}^{(2)}T = \frac{1}{3} \vartheta^\alpha \wedge T, \quad {}^{(2)}Q_{\alpha\beta} = 0, \quad (87)$$

are fulfilled, we derive in Appendix E in (E13) that $[\mathcal{R}ica := \vartheta^\alpha \wedge (e_\beta] {}^{(5)}W_\alpha{}^\beta)]$

$$\mathcal{R}ica - \frac{1}{3} d(2T + 3Q - \Lambda) = 0. \quad (88)$$

This equation can be understood as an integrability condition for (86). Substitution of (86) into (88) yields

$$0 = \mathcal{R}ica + \frac{1}{3} d\{(2\xi_0 - 1)\Lambda + 3[1 - 2(\xi_0 + \xi_1)]Q\}. \quad (89)$$

We have two free parameters. Thus, on this level, we can always find solutions with $\mathcal{R}ica = 0$. However, Eq.(89) also implies the non-trivial result

$$d\mathcal{R}ica = 0. \quad (90)$$

The second Bianchi identity may yield more conditions.

We now turn to the spherically symmetric solution of Sec.VII. In this case, we have the prolongation ansatz (84) with

$$\xi_0 = \frac{1}{2}, \quad \xi_1 = 0, \quad {}^{(3)}T^\alpha = 0. \quad (91)$$

Then (86) becomes

$$\Lambda - 3Q - 2T = 0. \quad (92)$$

This is consistent with the first Bianchi identity, see (E14). Since only ${}^{(2)}T^\alpha \neq 0$, we have

$$\Lambda - 3Q - 2T = \text{exact form}. \quad (93)$$

The distortion 1-form (10) can be taken from (66) and (69) or from [7], Eq.(35):

$$N_{\alpha\beta} = \frac{1}{2} Q_{\alpha\beta}. \quad (94)$$

Note that $N_{[\alpha\beta]} = 0$. In this special case, the curvature can be easily decomposed in Riemannian and post-Riemannian pieces,

$$R_\alpha{}^\beta = \tilde{R}_\alpha{}^\beta + \frac{1}{2} \tilde{D}Q_\alpha{}^\beta - \frac{1}{4} \mathcal{Q}_\alpha{}^\gamma \wedge \mathcal{Q}_\gamma{}^\beta, \quad (95)$$

where $\tilde{R}_\alpha{}^\beta$ denotes the purely Riemannian part of the curvature and $\tilde{D}$ the exterior covariant derivative with respect to the Riemannian connection.

Possibly, for a real “liberated” aether dynamics, one is forced to allow for $\mathcal{R}ica \neq 0$. Apparently $\mathcal{R}ica$ is the non-exact piece of Λ, Q, T and as such contributes generically to ${}^{(3)}Z_{\alpha\beta}$ and ${}^{(4)}Z_{\alpha\beta}$.

There is a further property of our specific exact solution which is of interest. The ansatz (84), together with the first Bianchi identity, yields

$$\begin{aligned} (Q_{\alpha\mu} \wedge Q_\beta{}^\mu + 4R_{[\alpha\beta]}) \wedge \vartheta^\beta &= \\ (\mathcal{Q}_{\alpha\mu} \wedge \mathcal{Q}_\beta{}^\mu + 4R_{[\alpha\beta]}) \wedge \vartheta^\beta &= 0. \end{aligned} \quad (96)$$

Compare now (95) with (96) and find

$$R_{\alpha\beta} \wedge \vartheta^\beta = \tilde{R}_{\alpha\beta} \wedge \vartheta^\beta + W_{\alpha\beta} \wedge \vartheta^\beta + \frac{1}{2} (\tilde{D}Q_{\alpha\beta}) \wedge \vartheta^\beta, \quad (97)$$

i.e., we have $\xi_0 = 1/2$ and $\xi_1 = 0$. This implies for the symmetric (strain) curvature

$$Z_{\alpha\beta} \wedge \vartheta^\beta = \tilde{R}_{\alpha\beta} \wedge \vartheta^\beta + \frac{1}{2} (\tilde{D}Q_{\alpha\beta}) \wedge \vartheta^\beta. \quad (98)$$

We decompose the curvature into symmetric and antisymmetric pieces. This yields

$$Z_{\alpha\beta} = \frac{1}{2} (\tilde{D}Q_{\alpha\beta}), \quad (99)$$

$$W_{\alpha\beta} = \tilde{R}_{\alpha\beta} - \frac{1}{4} \mathcal{Q}_{[\alpha}{}^\gamma \wedge \mathcal{Q}_{\beta]\gamma}. \quad (100)$$

Thus an extra field equation for $\mathcal{Q}_{\alpha\beta}$ is implied,

$$(\tilde{D}\mathcal{Q}_{\alpha\beta}) \wedge \vartheta^\beta = 0. \quad (101)$$

Generally, this equation implies further integrability conditions. With

$$DD\mathcal{Q}_{\alpha\beta} = -2R_{(\alpha}{}^\gamma \wedge \mathcal{Q}_{\beta)\gamma} \quad (102)$$

and the ansatz (84), we find the algebraic constraint

$$\frac{1}{\xi_0} \tilde{R}_{\alpha\beta} \wedge T^\beta = 0, \quad (103)$$

which has to be fulfilled by this solution.

X. DISCUSSION

We constructed a model within MAG which exhibits vector-like Lorentz violating fields. It may be seen as analogue to the Einstein-aether theory of Jacobson et al. We were able to find a simple exact spherically symmetric solution for the field equations of a truncated Lagrangian. Our solution distorts the conventional spherically symmetric Schwarzschild-de Sitter spacetime by nonmetricity and torsion. The presence of nonmetricity will obstruct the local Lorentz invariance of the Riemannian spacetime. Further investigations should include the search for wave-like aether solution which, most likely, will require a more complicated “background” than simple Schwarzschild spacetime, cf. plane-wave solutions in MAG [22, 50, 58].

It should be understood that MAG is a comprehensive framework for classical gravitational field theories. Different authors started with MAG and, by using the nonlinear realization technique, tried to “freeze out” certain degrees of freedom like, e.g., the nonmetricity. Percacci [62], Tresguerres and Mielke [79] (see also [38, 47]), and, most recently, Kirsch [40] developed models of such a kind, for earlier work one should compare Lord and Goswami [48, 49], see also Tresguerres [78]. There is not much doubt that gravity has a metric-affine structure. Therefore, MAG seems an appropriate framework for classical gravity. But there are different ways to realize this structure. Still, if Lorentz invariance turns out to be violated, then the nonmetricity of spacetime should play a leading role.

Acknowledgments

We thank Robert Bluhm, Ted Jacobson, Dirk Puetzfeld, Yakov Itin, Ingo Kirsch, and Romualdo Tresguerres for valuable comments on the manuscript. This project has been supported by the grant HE 528/20-1 of the DFG, Bonn.

APPENDIX A: IRREDUCIBLE DECOMPOSITION OF THE STRAIN

CURVATURE $Z_{\alpha\beta}$ IN n DIMENSIONS

The 2-form $Z_{\alpha\beta}$ in (32) — for n dimensions we have $1/n$ instead of $1/4$ — can be cut into different pieces by contraction with e_α , transvecting with ϑ^α , and by “hodge”-ing the corresponding expressions:

$$Z_\alpha := e^\beta \rfloor Z_{\alpha\beta}, \quad \hat{\Delta} := \frac{1}{n-2} \vartheta^\alpha \wedge Z_\alpha, \quad Y_\alpha := *(Z_{\alpha\beta} \wedge \vartheta^\beta). \quad (\text{A1})$$

Subsequently we can subtract out traces:

$$\Xi_\alpha := Z_\alpha - \frac{1}{2} e_\alpha \rfloor (\vartheta^\gamma \wedge Z_\gamma), \quad \Upsilon_\alpha := Y_\alpha - \frac{1}{n-2} e_\alpha \rfloor (\vartheta^\gamma \wedge Y_\gamma). \quad (\text{A2})$$

The irreducible pieces may then be written as (see [30])

$${}^{(2)}Z_{\alpha\beta} := -\frac{1}{2} *(\vartheta_{(\alpha} \wedge \Upsilon_{\beta)}), \quad (\text{A3})$$

$${}^{(3)}Z_{\alpha\beta} := \frac{1}{n+2} (n \vartheta_{(\alpha} \wedge e_{\beta)} \rfloor - 2 g_{\alpha\beta}) \hat{\Delta}, \quad (\text{A4})$$

$${}^{(4)}Z_{\alpha\beta} := \frac{1}{n} g_{\alpha\beta} Z, \quad (\text{A5})$$

$${}^{(5)}Z_{\alpha\beta} := \frac{2}{n} \vartheta_{(\alpha} \wedge \Xi_{\beta)}, \quad (\text{A6})$$

$${}^{(1)}Z_{\alpha\beta} := Z_{\alpha\beta} - {}^{(2)}Z_{\alpha\beta} - {}^{(3)}Z_{\alpha\beta} - {}^{(4)}Z_{\alpha\beta} - {}^{(5)}Z_{\alpha\beta}. \quad (\text{A7})$$

APPENDIX B: EXPRESSING THE CURVATURE ${}^{(3)}Z_{\alpha\beta}$ IN TERMS OF NONMETRICITY AND TORSION

According to (A4), the curvature ${}^{(3)}Z_{\alpha\beta}$ can be expressed in terms of $\hat{\Delta}$. Thus, we start from the definition of $\hat{\Delta}$ in (A1)₂ and move the interior product to the left:

$$\hat{\Delta} = \frac{1}{n-2} \vartheta^\alpha \wedge (e^\beta \rfloor Z_{\alpha\beta}) = \frac{1}{n-2} [-e^\beta \rfloor (\vartheta^\alpha \wedge Z_{\alpha\beta}) + Z_\alpha{}^\alpha] = -\frac{1}{n-2} e^\alpha \rfloor (Z_{\alpha\beta} \wedge \vartheta^\beta). \quad (\text{B1})$$

Obviously, we have to express $\mathbb{Z}_{\alpha\beta} \wedge \vartheta^\beta$ in terms of nonmetricity and torsion. This should be possible by means of the zeroth Bianchi identity

$$DQ_{\alpha\beta} = -DDg_{\alpha\beta} = R_{\alpha}{}^{\gamma} g_{\gamma\beta} + R_{\beta}{}^{\gamma} g_{\alpha\gamma} = 2R_{(\alpha\beta)} = 2Z_{\alpha\beta}. \quad (\text{B2})$$

We wedge with ϑ^β from the right and obtain

$$D(Q_{\alpha\beta} \wedge \vartheta^\beta) = 2Z_{\alpha\beta} \wedge \vartheta^\beta - Q_{\alpha\beta} \wedge T^\beta. \quad (\text{B3})$$

On the other hand, by making use of (25) (suitably generalized for n dimensions) and of

$$D\vartheta_\alpha = D(g_{\alpha\beta} \vartheta^\beta) = (Dg_{\alpha\beta}) \wedge \vartheta^\beta + g_{\alpha\beta} D\vartheta^\beta = -Q_{\alpha\beta} \wedge \vartheta^\beta + T_\alpha, \quad (\text{B4})$$

we can calculate

$$\begin{aligned} D(Q_{\alpha\beta} \wedge \vartheta^\beta) &\stackrel{(25)}{=} D\left(Q \wedge \vartheta_\alpha + \frac{1}{n-1} \vartheta_\alpha \wedge \Lambda + P_\alpha\right) \\ &= dQ \wedge \vartheta_\alpha - Q \wedge D\vartheta_\alpha + \frac{1}{n-1} D\vartheta_\alpha \wedge \Lambda - \frac{1}{n-1} \vartheta_\alpha \wedge d\Lambda + DP_\alpha \\ &\stackrel{(\text{B4})}{=} dQ \wedge \vartheta_\alpha - Q \wedge (-Q_{\alpha\beta} \wedge \vartheta^\beta + T_\alpha) \\ &\quad + \frac{1}{n-1} (-Q_{\alpha\beta} \wedge \vartheta^\beta + T_\alpha) \wedge \Lambda - \frac{1}{n-1} \vartheta_\alpha \wedge d\Lambda + DP_\alpha \\ &\stackrel{(25)}{=} dQ \wedge \vartheta_\alpha + Q \wedge \left(Q \wedge \vartheta_\alpha + \frac{1}{n-1} \vartheta_\alpha \wedge \Lambda + P_\alpha\right) - Q \wedge T_\alpha \\ &\quad - \frac{1}{n-1} \left(Q \wedge \vartheta_\alpha + \frac{1}{n-1} \vartheta_\alpha \wedge \Lambda + P_\alpha\right) \wedge \Lambda + \frac{1}{n-1} T_\alpha \wedge \Lambda \\ &\quad - \frac{1}{n-1} \vartheta_\alpha \wedge d\Lambda + DP_\alpha \\ &= dQ \wedge \vartheta_\alpha + Q \wedge Q \wedge \vartheta_\alpha + \frac{1}{n-1} Q \wedge \vartheta_\alpha \wedge \Lambda + Q \wedge P_\alpha - Q \wedge T_\alpha \\ &\quad - \frac{1}{n-1} Q \wedge \vartheta_\alpha \wedge \Lambda - \frac{1}{(n-1)^2} \vartheta_\alpha \wedge \Lambda \wedge \Lambda - \frac{1}{n-1} P_\alpha \wedge \Lambda + \frac{1}{n-1} T_\alpha \wedge \Lambda \\ &\quad - \frac{1}{n-1} \vartheta_\alpha \wedge d\Lambda + DP_\alpha \\ &= dQ \wedge \vartheta_\alpha + Q \wedge P_\alpha - Q \wedge T_\alpha - \frac{1}{n-1} P_\alpha \wedge \Lambda + \frac{1}{n-1} T_\alpha \wedge \Lambda \\ &\quad - \frac{1}{n-1} \vartheta_\alpha \wedge d\Lambda + DP_\alpha. \end{aligned} \quad (\text{B5})$$

Now we can compare (B3) and (B5). We find

$$\begin{aligned} 2 \mathbb{Z}_{\alpha\beta} \wedge \vartheta^\beta &= 2Z_{\alpha\beta} \wedge \vartheta^\beta - 2^{(4)} Z_{\alpha\beta} \wedge \vartheta^\beta \\ &= Q \wedge P_\alpha - Q \wedge T_\alpha - \frac{1}{n-1} P_\alpha \wedge \Lambda \\ &\quad + \frac{1}{n-1} T_\alpha \wedge \Lambda - \frac{1}{n-1} \vartheta_\alpha \wedge d\Lambda + DP_\alpha + Q_{\alpha\beta} \wedge T^\beta. \end{aligned} \quad (\text{B6})$$

We expand the last term by means of the irreducible decomposition of torsion and non-metricity:

$$\begin{aligned}
Q_{\alpha\beta} \wedge T^\beta &= \mathcal{Q}_{\alpha\beta} \wedge T^\beta + Q \wedge T_\alpha \\
&= \mathcal{Q}_{\alpha\beta} \wedge \left({}^{(1)}T^\beta + {}^{(3)}T^\beta \right) + \mathcal{Q}_{\alpha\beta} \wedge \left(\frac{1}{n-1} \vartheta^\beta \wedge T \right) + Q \wedge T_\alpha \\
&= \mathcal{Q}_{\alpha\beta} \wedge \left({}^{(1)}T^\beta + {}^{(3)}T^\beta \right) + \frac{1}{(n-1)^2} \vartheta_\alpha \wedge \Lambda \wedge T + \frac{1}{n-1} P_\alpha \wedge T + Q \wedge T_\alpha. \quad (\text{B7})
\end{aligned}$$

By the same token,

$$\frac{1}{n-1} T_\alpha \wedge \Lambda = \frac{1}{n-1} \left({}^{(1)}T_\alpha + {}^{(3)}T_\alpha \right) \wedge \Lambda + \frac{1}{(n-1)^2} \vartheta_\alpha \wedge T \wedge \Lambda. \quad (\text{B8})$$

Substituting (B7,B8) into (B6) yields

$$\begin{aligned}
2 \mathcal{Z}_{\alpha\beta} \wedge \vartheta^\beta &= DP_\alpha - \frac{1}{n-1} \vartheta_\alpha \wedge d\Lambda + \mathcal{Q}_{\alpha\beta} \wedge \left({}^{(1)}T^\beta + {}^{(3)}T^\beta \right) \\
&\quad + \frac{1}{n-1} \left({}^{(1)}T_\alpha + {}^{(3)}T_\alpha \right) \wedge \Lambda + P_\alpha \wedge \left[Q - \frac{1}{n-1} (\Lambda - T) \right]. \quad (\text{B9})
\end{aligned}$$

We use the following properties of the irreducible pieces:

$$e^\alpha \rfloor P_\alpha = e^\alpha \rfloor {}^{(1)}T_\alpha = e^\alpha \rfloor {}^{(3)}T_\alpha = 0, \quad e^\alpha \rfloor \mathcal{Q}_{\alpha\beta} = \Lambda_\beta. \quad (\text{B10})$$

Then we find

$$\begin{aligned}
2e^\alpha \rfloor (\mathcal{Z}_{\alpha\beta} \wedge \vartheta^\beta) &= P^\alpha e_\alpha \rfloor \left[Q - \frac{1}{n-1} (\Lambda - T) \right] + \frac{n}{n-1} \left({}^{(1)}T_\alpha + {}^{(3)}T_\alpha \right) \wedge \Lambda^\alpha \\
&\quad - \frac{n-2}{n-1} d\Lambda + e^\alpha \rfloor DP_\alpha - \mathcal{Q}_{\alpha\beta} \wedge e^\alpha \rfloor \left({}^{(1)}T^\beta + {}^{(3)}T^\beta \right). \quad (\text{B11})
\end{aligned}$$

We can further simplify the last term. First we note that

$$e^\alpha \rfloor {}^{(3)}T^\beta = (-1)^s e^\alpha \rfloor \frac{1}{3} \star [\vartheta^\beta \wedge \star (T^\gamma \wedge \vartheta_\gamma)] = (-1)^s \frac{1}{3} \star [\vartheta^\beta \wedge \star (T^\gamma \wedge \vartheta_\gamma) \wedge \vartheta^\alpha] = -e^\beta \rfloor {}^{(3)}T^\alpha. \quad (\text{B12})$$

Hence,

$$\begin{aligned}
\mathcal{Q}_{\alpha\beta} \wedge e^\alpha \rfloor \left({}^{(1)}T^\beta + {}^{(3)}T^\beta \right) &= \mathcal{Q}_{\alpha\beta} \wedge e^\alpha \rfloor {}^{(1)}T^\beta = {}^{(1)}Q_{\alpha\beta} \wedge e^\alpha \rfloor {}^{(1)}T^\beta \\
&\quad + {}^{(2)}Q_{\alpha\beta} \wedge e^\alpha \rfloor {}^{(1)}T^\beta + {}^{(3)}Q_{\alpha\beta} \wedge e^\alpha \rfloor {}^{(1)}T^\beta. \quad (\text{B13})
\end{aligned}$$

The last term can be further rewritten as

$${}^{(3)}Q_{\alpha\beta} \wedge e^\alpha \rfloor {}^{(1)}T^\beta = \frac{2n}{(n-1)(n+2)} \left(\vartheta_{(\alpha} \Lambda_{\beta)} - \frac{1}{n} g_{\alpha\beta} \Lambda \right) \wedge e^\alpha \rfloor {}^{(1)}T^\beta$$

$$\begin{aligned}
&= \frac{n}{(n-1)(n+2)} (\vartheta_\beta \Lambda_\alpha \wedge e^\alpha \rfloor^{(1)} T^\beta + \Lambda_\beta \vartheta_\alpha e^\alpha \rfloor^{(1)} T^\beta) \\
&= \frac{n}{(n-1)(n+2)} [\Lambda_\alpha (-e^\alpha \rfloor (\vartheta_\beta \wedge^{(1)} T^\beta) + {}^{(1)} T^\alpha) + 2\Lambda_\beta {}^{(1)} T^\beta] \\
&= \frac{3n}{(n-1)(n+2)} \Lambda_\alpha {}^{(1)} T^\alpha, \tag{B14}
\end{aligned}$$

where we used $\vartheta_\beta \wedge^{(1)} T^\beta = 0$. Finally we arrive at

$$\begin{aligned}
\hat{\Delta} &= \frac{1}{2(n-1)} d\Lambda - \frac{1}{2(n-2)} e^\alpha \rfloor DP_\alpha \\
&\quad - \frac{1}{2(n-2)} \left\{ \frac{1}{n-1} P_\alpha e^\alpha \rfloor [(n-1)Q + \Lambda - T] + \left(\frac{n+1}{n+2} {}^{(1)} T_\alpha + \frac{n}{n-1} {}^{(3)} T_\alpha \right) \Lambda^\alpha \right. \\
&\quad \left. - ({}^{(1)} Q_{\alpha\beta} + {}^{(2)} Q_{\alpha\beta}) \wedge e^\alpha \rfloor^{(1)} T^\beta \right\}. \tag{B15}
\end{aligned}$$

Note that in the last line we could substitute ${}^{(2)} Q_{\alpha\beta} = -2 e_{(\alpha} \rfloor P_{\beta)}/3$.

APPENDIX C: THE ${}^{(3)} Z^{\alpha\beta} \wedge {}^\star({}^{(3)} Z_{\alpha\beta})$ PIECE OF THE LAGRANGIAN

We proof that the following relation holds for *arbitrary* spacetimes:

$${}^{(3)} Z^{\alpha\beta} \wedge {}^\star({}^{(3)} Z_{\alpha\beta}) = \frac{n(n-2)}{n+2} \hat{\Delta} \wedge {}^\star \hat{\Delta}. \tag{C1}$$

This comes about since ${}^{(3)} Z^{\alpha\beta}$ corresponds to a *scalar*-valued degree of freedom, namely to the two-form $\hat{\Delta}$, see (A4). For a p-form ϕ , we have the rules for the Hodge dual ${}^\star\phi = (-1)^{p(n-p)-1}\phi$ in the case of Lorentz signature, furthermore, $\vartheta^\alpha \wedge (e_\alpha \rfloor \phi) = p\phi$ and ${}^\star(\phi \wedge \vartheta_\alpha) = e_\alpha \rfloor {}^\star\phi$. Thus, the terms quadratic in contractions of $\hat{\Delta}$ in the end evaluate to $\hat{\Delta} \wedge {}^\star \hat{\Delta}$,

$$(e^\alpha \rfloor \hat{\Delta}) \wedge {}^\star(e_\alpha \rfloor \hat{\Delta}) = -(e^\alpha \rfloor \hat{\Delta}) \wedge {}^\star(e_\alpha \rfloor {}^\star \hat{\Delta}) = -(e^\alpha \rfloor \hat{\Delta}) \wedge \vartheta_\alpha \wedge {}^\star \hat{\Delta} = 2 \hat{\Delta} \wedge {}^\star \hat{\Delta}. \tag{C2}$$

We recall the definition (A4)

$${}^{(3)} Z_{\alpha\beta} = \frac{1}{n+2} \left[n \vartheta_{(\alpha} \wedge (e_{\beta)} \rfloor \hat{\Delta} - 2 g_{\alpha\beta} \hat{\Delta} \right]. \tag{C3}$$

It is symmetric ${}^{(3)} Z^{[\alpha\beta]} = 0$ and tracefree ${}^{(3)} Z^\gamma{}_\gamma = 0$. Thus,

$$\begin{aligned}
{}^{(3)} Z_{\alpha\beta} \wedge {}^\star({}^{(3)} Z^{\alpha\beta}) &= \frac{1}{n+2} \left[n \vartheta_\alpha \wedge (e_\beta \rfloor \hat{\Delta}) \wedge {}^\star({}^{(3)} Z^{\alpha\beta}) \right] \\
&= \frac{1}{(n+2)^2} \left[n^2 \vartheta_\alpha \wedge (e_\beta \rfloor \hat{\Delta}) \wedge {}^\star \left(\vartheta^{(\alpha} \wedge (e^{\beta)} \rfloor \hat{\Delta}) \right) - 2n \vartheta_\alpha \wedge (e_\beta \rfloor \hat{\Delta}) \wedge g^{\alpha\beta} {}^\star \hat{\Delta} \right]. \tag{C4}
\end{aligned}$$

In order to calculate the first term, we apply the rules for commuting the Hodge star with the exterior/interior product.

$$\begin{aligned}
\star \left(\vartheta^{(\alpha} \wedge (e^{\beta)} \rfloor \hat{\Delta} \right) &= -\star \left[(e^{(\beta)} \rfloor \hat{\Delta}) \wedge \vartheta^{\alpha} \right] = -e^{(\alpha} \rfloor \star (e^{\beta)} \rfloor \hat{\Delta}) \\
&= -(-1)^{n-1} e^{(\alpha} \rfloor \left(\star \hat{\Delta} \wedge \vartheta^{\beta} \right) = e^{(\alpha} \rfloor \left(\vartheta^{\beta} \wedge \star \hat{\Delta} \right) \\
&= g^{\alpha\beta} \star \hat{\Delta} - \vartheta^{(\alpha} \wedge (e^{\beta)} \rfloor \star \hat{\Delta}).
\end{aligned} \tag{C5}$$

The second term in the brackets of (C4) simply evaluates to

$$-2n \vartheta_{\alpha} \wedge (e_{\beta} \rfloor \hat{\Delta}) \wedge g^{\alpha\beta} \star \hat{\Delta} = -2n \vartheta^{\alpha} \wedge (e_{\alpha} \rfloor \hat{\Delta}) \wedge \star \hat{\Delta} = -4n \hat{\Delta} \wedge \star \hat{\Delta}. \tag{C6}$$

Substituting (C5) and (C6) into (C4), we find

$${}^{(3)}Z^{\alpha\beta} \wedge \star {}^{(3)}Z^{\alpha\beta} = \frac{1}{(n+2)^2} \left[2n^2 \hat{\Delta} \wedge \star \hat{\Delta} - n^2 \vartheta_{\alpha} \wedge (e_{\beta} \rfloor \hat{\Delta}) \wedge \vartheta^{(\alpha} \wedge (e^{\beta)} \rfloor \star \hat{\Delta}) - 4n \hat{\Delta} \wedge \star \hat{\Delta} \right]. \tag{C7}$$

The term in the middle yields ($\vartheta^{\alpha} \wedge \vartheta_{\alpha} = 0$)

$$\begin{aligned}
(\vartheta_{\alpha} \wedge e_{\beta} \rfloor \hat{\Delta}) \wedge (\vartheta^{(\alpha} \wedge e^{\beta)} \rfloor \star \hat{\Delta}) &= \frac{1}{2} \left[\vartheta_{\alpha} \wedge (e_{\beta} \rfloor \hat{\Delta}) \wedge \vartheta^{\alpha} \wedge (e^{\beta} \rfloor \star \hat{\Delta}) + \vartheta_{\alpha} \wedge (e_{\beta} \rfloor \hat{\Delta}) \wedge \vartheta^{\beta} \wedge (e^{\alpha} \rfloor \star \hat{\Delta}) \right] \\
&= \frac{1}{2} \left[-\vartheta^{\beta} \wedge (e_{\beta} \rfloor \hat{\Delta}) \wedge \vartheta_{\alpha} \wedge (e^{\alpha} \rfloor \star \hat{\Delta}) \right] = -(n-2) \hat{\Delta} \wedge \star \hat{\Delta}.
\end{aligned} \tag{C8}$$

Eventually,

$${}^{(3)}Z^{\alpha\beta} \wedge \star {}^{(3)}Z^{\alpha\beta} = \frac{1}{(n+2)^2} \left[2n^2 + n^2(n-2) - 4n \right] \hat{\Delta} \wedge \star \hat{\Delta} = \frac{n(n-2)}{n+2} \hat{\Delta} \wedge \star \hat{\Delta}. \tag{C9}$$

APPENDIX D: THE MAG LAGRANGIAN QUADRATIC IN CURVATURE, TORSION, AND NONMETRICITY

The most general parity conserving quadratic Lagrangian, which is expressed in terms of the $4 + 3 + 6 + 5$ irreducible pieces of $Q_{\alpha\beta}$, T^{α} , $W_{\alpha}{}^{\beta}$, and $Z_{\alpha}{}^{\beta}$, respectively, reads (see [19], [59], [29], and references given):

$$\begin{aligned}
V_{\text{MAG}} &= \frac{1}{2\kappa} \left[-a_0 R^{\alpha\beta} \wedge \eta_{\alpha\beta} - 2\lambda_0 \eta + T^{\alpha} \wedge \star \left(\sum_{I=1}^3 a_I {}^{(I)}T_{\alpha} \right) \right. \\
&\quad + 2 \left(\sum_{I=2}^4 c_I {}^{(I)}Q_{\alpha\beta} \right) \wedge \vartheta^{\alpha} \wedge \star T^{\beta} + Q_{\alpha\beta} \wedge \star \left(\sum_{I=1}^4 b_I {}^{(I)}Q^{\alpha\beta} \right) \\
&\quad \left. + b_5 ({}^{(3)}Q_{\alpha\gamma} \wedge \vartheta^{\alpha}) \wedge \star ({}^{(4)}Q^{\beta\gamma} \wedge \vartheta_{\beta}) \right]
\end{aligned}$$

$$\begin{aligned}
& - \frac{1}{2\rho} R^{\alpha\beta} \wedge * \left(\sum_{I=1}^6 w_I {}^{(I)}W_{\alpha\beta} + w_7 \vartheta_\alpha \wedge (e_\gamma \rfloor {}^{(5)}W^\gamma{}_\beta) \right. \\
& \left. + \sum_{I=1}^5 z_I {}^{(I)}Z_{\alpha\beta} + z_6 \vartheta_\gamma \wedge (e_\alpha \rfloor {}^{(2)}Z^\gamma{}_\beta) + \sum_{I=7}^9 z_I \vartheta_\alpha \wedge (e_\gamma \rfloor {}^{(I-4)}Z^\gamma{}_\beta) \right) \quad . \quad (D1)
\end{aligned}$$

Here κ is the dimensionful (weak) gravitational constant, λ_0 the “bare” cosmological constant, and the dimensionless ρ is the strong gravity coupling constant. The constants $a_0, \dots, a_3, b_1, \dots, b_5, c_2, c_3, c_4, w_1, \dots, w_7, z_1, \dots, z_9$ are dimensionless and should be of order unity. Note the nontrivial formula for the Hilbert-Einstein type of Lagrangian

$$R_{\alpha\beta} \wedge \eta^{\alpha\beta} = {}^{(6)}W_{\alpha\beta} \wedge \eta^{\alpha\beta} . \quad (D2)$$

Because of the irreducibility, we have

$$\begin{aligned}
T^2 & \sim T^\alpha \wedge * (a_1 {}^{(1)}T_\alpha + a_2 {}^{(2)}T_\alpha + a_3 {}^{(3)}T_\alpha) \\
& = a_1 {}^{*(1)}T_\alpha \wedge {}^{(1)}T_\alpha + a_2 {}^{*(2)}T_\alpha \wedge {}^{(2)}T_\alpha + a_3 {}^{*(3)}T_\alpha \wedge {}^{(3)}T_\alpha , \quad (D3)
\end{aligned}$$

and similar formulas for the pure square pieces of $Q_{\alpha\beta}$, $W_{\alpha\beta}$, and $Z_{\alpha\beta}$. In the curvature square terms in (D1) we introduced the irreducible pieces of the antisymmetric part $W_{\alpha\beta} := R_{[\alpha\beta]}$ (rotation part) and the symmetric part $Z_{\alpha\beta} := R_{(\alpha\beta)}$ (strain part) of the curvature 2-form $R_{\alpha\beta}$. In $Z_{\alpha\beta}$, we have the purely *post*-Riemannian part of the curvature. Note the peculiar cross terms with c_I and b_5 , and the *exotic* terms with w_7 and z_6, z_7, z_8, z_9 .

In the component formalism, Esser [19] has carefully enumerated all different pieces of the quadratic MAG Lagrangian; for the corresponding nonmetricity and torsion pieces, see also Duan et al. [15]. Accordingly, Eq.(D1) represents the most general quadratic parity-conserving MAG-Lagrangian. All previously published quadratic parity-conserving Lagrangians are subcases of (D1). Hence (D1) is a safe starting point for our future considerations.

APPENDIX E: THE 2-FORM *Rica* AND THE FIRST BIANCHI IDENTITY

The first Bianchi identity $B^\alpha \equiv 0$, with the 3-form

$$B^\alpha := DT^\alpha - R_\beta{}^\alpha \wedge \vartheta^\beta , \quad (E1)$$

interrelates torsion and curvature. The irreducible decomposition of the first Bianchi identity reads (see [30])

$${}^{(2)}B^\alpha = \frac{1}{n-2} (e_\beta \rfloor B^\beta) \wedge \vartheta^\alpha, \quad (\text{E2})$$

$${}^{(3)}B^\alpha = \frac{1}{4} e_\alpha \rfloor (\vartheta^\beta \wedge B_\beta), \quad (\text{E3})$$

$${}^{(1)}B^\alpha = B^\alpha - {}^{(2)}B^\alpha - {}^{(3)}B^\alpha. \quad (\text{E4})$$

In the last term of (E1), because of the contraction with the coframe, some of the irreducible components of the curvature drop out, see [30], Eqs.(B.4.15) and (B.4.29):

$$R_\beta{}^\alpha \wedge \vartheta^\beta = ({}^{(2)}W_\beta{}^\alpha + {}^{(3)}W_\beta{}^\alpha + {}^{(5)}W_\beta{}^\alpha + {}^{(2)}Z_\beta{}^\alpha + {}^{(3)}Z_\beta{}^\alpha + {}^{(4)}Z_\beta{}^\alpha) \wedge \vartheta^\beta. \quad (\text{E5})$$

We contract this piece by the frame:

$$\begin{aligned} e_\alpha \rfloor (R_\beta{}^\alpha \wedge \vartheta^\beta) &= e_\alpha \rfloor ({}^{(5)}W_\beta{}^\alpha \wedge \vartheta^\beta) + e_\alpha \rfloor ({}^{(3)}Z_\beta{}^\alpha \wedge \vartheta^\beta) + e_\alpha \rfloor ({}^{(5)}Z_\beta{}^\alpha \wedge \vartheta^\beta) \\ &= -\mathcal{R}ica - (n-2) \hat{\Delta} + \frac{n-2}{2} dQ. \end{aligned} \quad (\text{E6})$$

We introduced as abbreviation the antisymmetric rotational Ricci 2-form for $W_{\alpha\beta}$ by

$$\mathcal{R}ica := \vartheta^\alpha \wedge (e_\beta \rfloor {}^{(5)}W_\alpha{}^\beta) = \vartheta^\alpha \wedge (e_\beta \rfloor W_\alpha{}^\beta), \quad (\text{E7})$$

see [30], Eqs.(B.4.8) and (B.4.11). $\mathcal{R}ica$ has $n(n-1)/2$ independent components. Note that there is a subtlety involved here. The (premetric) Ricci 1-form is usually defined in terms of the *total* curvature according to $\text{Ric}_\alpha := e_\beta \rfloor R_\alpha{}^\beta$. Here, however, our definition refers only to the rotational curvature $W_{\alpha\beta}$, see (29). Similarly we obtain

$$\vartheta^\alpha \wedge (R_{\beta\alpha} \wedge \vartheta^\beta) = \vartheta^\alpha \wedge ({}^{(3)}W_{\beta\alpha} \wedge \vartheta^\beta) =: \hat{X}. \quad (\text{E8})$$

Before we substitute (E6) and (E8) into the first Bianchi identity, we should first compute the corresponding torsion pieces:

$$\begin{aligned} e_\alpha \rfloor DT^\alpha &= e_\alpha \rfloor [D({}^{(1)}T^\alpha + {}^{(3)}T^\alpha)] + \frac{1}{n-1} ({}^{(1)}T^\alpha + {}^{(3)}T^\alpha) e_\alpha \rfloor T \\ &\quad - \frac{n-2}{n-1} dT, \end{aligned} \quad (\text{E9})$$

$$\vartheta^\alpha \wedge DT_\alpha = -D(\vartheta^\alpha \wedge {}^{(3)}T_\alpha) + T^\alpha \wedge T_\alpha. \quad (\text{E10})$$

In this way, we find for the 3-form B^α eventually,

$$\begin{aligned} e_\beta \rfloor B^\beta &= e_\alpha \rfloor [D({}^{(1)}T^\alpha + {}^{(3)}T^\alpha)] + \frac{1}{n-1} ({}^{(1)}T^\alpha + {}^{(3)}T^\alpha) e_\alpha \rfloor T \\ &\quad - \frac{n-2}{n-1} dT + \mathcal{R}ica + (n-2) \hat{\Delta} - \frac{n-2}{2} dQ = 0, \end{aligned} \quad (\text{E11})$$

$$\vartheta^\beta \wedge B_\beta = -D(\vartheta^\alpha \wedge {}^{(3)}T_\alpha) + T^\alpha \wedge T_\alpha - \hat{X} = 0. \quad (\text{E12})$$

Provided the torsion possesses *only its trace piece*, that is $^{(1)}T^\alpha = ^{(3)}T^\alpha = 0$, and furthermore $^{(2)}Q_{\alpha\beta} = 0$, see the constraints (38), we have $\hat{\Delta} = d\Lambda/[2(n-1)]$ and

$$\mathcal{R}ica - \frac{n-2}{2(n-1)} d[2T + (n-1)Q - \Lambda] = 0. \quad (\text{E13})$$

Thus, in $n = 4$, with the constraints (38) fulfilled and for $\mathcal{R}ica = 0$, the first Bianchi identity yields

$$d(3Q + 2T - \Lambda) = 0. \quad (\text{E14})$$

Accordingly, under these conditions, the Weyl, the torsion, and the shear 1-forms are algebraically related to each other.

APPENDIX F: CURVATURE OF THE SPHERICALLY SYMMETRIC AETHER SOLUTION OF SEC.VII

With the help of our Reduce-Excalc computer algebra programs, we calculate the rotational and the strain curvature, respectively (the diamonds $\diamond$ and the bullets $\bullet$ denote those matrix elements that are already known because of the antisymmetry or the symmetry of the matrix involved):

$$^{(1)}W_{\alpha\beta} = \left(\frac{m}{r^3} - \frac{(\ell_0 + \ell_1)(4\ell_1 - \ell_0)}{96r^4 e^{2\mu(r)}} \right) \begin{pmatrix} 0 & 2\vartheta^{01} & -\vartheta^{02} & -\vartheta^{03} \\ \diamond & 0 & \vartheta^{12} & \vartheta^{13} \\ \diamond & \diamond & 0 & -2\vartheta^{23} \\ \diamond & \diamond & \diamond & 0 \end{pmatrix} \quad (\text{weyl}), \quad (\text{F1})$$

$$^{(2)}W_{\alpha\beta} = 0 \quad (\text{paircom} = 0), \quad (\text{F2})$$

$$^{(3)}W_{\alpha\beta} = 0 \quad (\text{pscalar} = 0), \quad (\text{F3})$$

$$^{(4)}W_{\alpha\beta} = -\vartheta_{[\alpha} \wedge \phi_{\beta]} \quad (\text{ricsymf}), \quad (\text{F4})$$

$$\begin{aligned} \phi_0 &= \frac{(\ell_0 + 7\ell_1)(\ell_0 - \ell_1)\vartheta^0 - 4\ell_0\ell_1\vartheta^1}{32r^4 e^{2\mu(r)}}, \\ \phi_1 &= -\frac{4\ell_0\ell_1\vartheta^0 + (\ell_0 - 5\ell_1)(\ell_0 - \ell_1)\vartheta^1}{32r^4 e^{2\mu(r)}}, \\ \phi_2 &= \frac{(\ell_0 + \ell_1)(\ell_0 - \ell_1)}{32r^4 e^{2\mu(r)}} \vartheta^2, \\ \phi_3 &= \frac{(\ell_0 + \ell_1)(\ell_0 - \ell_1)}{32r^4 e^{2\mu(r)}} \vartheta^3, \end{aligned}$$

$$^{(5)}W_{\alpha\beta} = 0 \quad (\text{ricanti} = 0), \quad (\text{F5})$$

$${}^{(6)}W_{\alpha\beta} = -\frac{1}{12} W \vartheta_{\alpha\beta}, \quad W = 4\lambda_0 + \frac{(\ell_0 + 5\ell_1)(\ell_0 + \ell_1)}{8r^4 e^{2\mu(r)}} \quad (\text{scalar}). \quad (\text{F6})$$

$${}^{(1)}Z_{\alpha\beta} = \frac{1}{48r^4 e^{2\mu(r)}} \begin{pmatrix} -4A_1 \vartheta^{01} & -2A_2 \vartheta^{01} & A_2 \vartheta^{02} + 2A_1 \vartheta^{12} & A_2 \vartheta^{03} + 2A_1 \vartheta^{13} \\ \bullet & 4A_3 \vartheta^{01} & -2A_3 \vartheta^{02} + A_2 \vartheta^{12} & -2A_3 \vartheta^{03} + A_2 \vartheta^{13} \\ \bullet & \bullet & 24\ell_1 r e^{2\mu(r)} \vartheta^{01} & 0 \\ \bullet & \bullet & \bullet & 24\ell_1 r e^{2\mu(r)} \vartheta^{01} \end{pmatrix}, \quad (\text{F7})$$

$$A_1 = \ell_1(4\lambda_0 r^3 + 15m - 9r), \quad A_2 = \ell_0(2\lambda_0 r^3 + 3m - 3r) + 9\ell_1(3m - r),$$

$$A_3 = 3\ell_1(3m - r),$$

$${}^{(2)}Z_{\alpha\beta} = 0, \quad (\text{F8})$$

$${}^{(3)}Z_{\alpha\beta} = 0, \quad (\text{F9})$$

$${}^{(4)}Z_{\alpha\beta} = -\frac{\ell_1}{2r^3} g_{\alpha\beta} \vartheta^{01} \quad (\text{dilcurv}), \quad (\text{F10})$$

$${}^{(5)}Z_{\alpha\beta} = \frac{1}{2} \vartheta_{(\alpha} \wedge \Xi_{\beta)}, \quad (\text{F11})$$

$$\Xi_0 = -\frac{(\lambda_0 r^3 - 3m)(\ell_0 + \ell_1) \vartheta^0 - 2(3m - r)\ell_1 \vartheta^1}{4r^4 e^{2\mu(r)}},$$

$$\Xi_1 = -\frac{[3(\ell_0 + 9\ell_1)m - 6(\ell_0 + 3\ell_1)r + (5\ell_0 + 9\ell_1)\lambda_0 r^3] \vartheta^1 - 6(3m - r)\ell_1 \vartheta^0}{12r^4 e^{2\mu(r)}},$$

$$\Xi_2 = -\frac{\ell_0 + 3\ell_1}{4r^3} \vartheta^2,$$

$$\Xi_3 = -\frac{\ell_0 + 3\ell_1}{4r^3} \vartheta^3.$$

- [1] M. Adak, T. Dereli and L. H. Ryder, *On the possible effects of spacetime non-metricity on neutrino oscillations*, Phys. Rev. **D69** (2004) 123002 (7 pages), [gr-qc/0303080](#).
- [2] R. E. Allen and S. Yokoo, *Searching for Lorentz violation*, Nucl. Phys. Proc. Suppl. **134**, 139–146 (2004), [hep-th/0402154](#).
- [3] A.A. Andrianov, R. Soldati and L. Sorbo, *Dynamical Lorentz symmetry breaking from 3+1 axion-Wess-Zumino model*, Phys. Rev. **D59** (1999) 025002 (13 pages), [hep-th/9806220](#).
- [4] O.V. Babourova and B.N. Frolov, *Matter with dilaton charge in Weyl-Cartan spacetime and evolution of the Universe*, Class. Quant. Grav. **20** (2003) 1423–1442, [gr-qc/0209077](#).
- [5] P. Baekler, *Prolongation in Metric-Affine Gravity*, unpublished manuscript (2003).

- [6] P. Baekler, R. Hecht, F.W. Hehl and T. Shirafuji, *Mass and spin of exact solutions of the Poincaré gauge theory*, Prog. Theor. Phys. **78** (1987) 16–21.
- [7] P. Baekler, F.W. Hehl, *Metric-affine gauge theory of gravity III. Prolongation and new classes of exact solutions*, in preparation (2005).
- [8] J.F. Barbero G. and E.J.S. Villaseñor, *Lorentz violations and Euclidean signature metrics*, Phys. Rev. **D68** (2003) 087501 (2 pages), [gr-qc/0307066](#).
- [9] M. Blagojević, *Gravitation and Gauge Symmetries* (IOP Publishing: Bristol, UK, 2002).
- [10] R. Bluhm, *Breaking Lorentz symmetry*, Phys. World **17N3** (2004) 41–46.
- [11] R. Bluhm and V.A. Kostelecky, *Spontaneous Lorentz violation, Nambu-Goldstone modes, and gravity*, [hep-th/0412320](#), 16 pages (2004).
- [12] S.M. Carroll and E.A. Lim, *Lorentz-violating vector fields slow the universe down*, Phys. Rev. **D70** (2004) 123525 (6 pages), [hep-th/0407149](#).
- [13] C.-M. Chen, J.M. Nester, R.-S. Tung, *Quasilocal energy momentum for gravity theories*, Phys. Lett. **A203** (1995) 5–11, [gr-qc/9411048](#).
- [14] T. Dereli, M. Önder, J. Schray, R.W. Tucker, and C. Wang, *Non-Riemannian gravity and the Einstein-Proca system*. Class. Quantum Grav. **13** (1996) L103–L109.
- [15] Y.-S. Duan, J.-C. Liu, X.-G. Dong, *General covariant energy-momentum conservation law in general spacetime*, Gen. Relat. Grav. **20** (1988) 485–496.
- [16] A. Einstein, Foreword in: M. Pantaleo, ed., *Cinquant’anni di Relatività 1905–1955* (Edizioni Giuntine and Sansoni Editore: Firenze, 1955), our translation from the German original.
- [17] C. Eling and T. Jacobson, *Static post-Newtonian equivalence of general relativity and gravity with a dynamical preferred frame*, Phys. Rev. **D69** (2004) 064005 (6 pages), [gr-qc/0310044](#).
- [18] C. Eling, T. Jacobson, D. Mattingly: *Einstein-Aether Theory*, 17 pages, to appear in the Proceedings of Deserfest (World Scientific), [gr-qc/0410001](#)
- [19] W. Esser, *Exact solutions of the metric-affine gauge theory of gravity*. Diploma Thesis, University of Cologne (1996).
- [20] R. Floreanini and R. Percacci, *The renormalization group flow of the dilaton potential*, Phys. Rev. **D52** (1995) 896–911, [hep-th/9412181](#).
- [21] B.Z. Foster, *Metric redefinitions in Einstein-aether theory*, 10 pages, [gr-qc/0502066](#).
- [22] A. García, A. Macías, D. Puetzfeld, and J. Socorro, *Plane-fronted waves in metric-affine gravity*, Phys. Rev. **D62** (2000) 044021 (7 pages), [gr-qc/0005038](#).

- [23] M. Gasperini, *Quasi-riemannian gravity and spontaneous breaking of the lorentz gauge symmetry in more than four-dimensions*, Phys. Rev. **D33** (1986) 3594–3606.
- [24] M. Gasperini, *Singularity prevention and broken Lorentz symmetry*, Class. Quantum Grav. **4** (1987) 485–494.
- [25] H. F. M. Goenner, *On the history of unified field theories*, Living Rev. Rel. **7** (2004) 2 (151 pages).
- [26] M.L. Graesser, A. Jenkins and M.B. Wise, *Spontaneous Lorentz violation and the long-range gravitational preferred-frame effect*, 10 pages, hep-th/0501223 (2005).
- [27] F. Gronwald and F.W. Hehl, *On the gauge aspects of gravity*, in: *International School of Cosmology and Gravitation: 14th Course: Quantum Gravity*, held May 1995 in Erice, Italy. Proceedings. P.G. Bergmann et al. (eds.). World Scientific, Singapore (1996) pp. 148–198, gr-qc/9602013.
- [28] M.P. Haugan, C. Lämmerzahl, *Principles of equivalence: their role in gravitation physics and experiments that test them*, Lect. Notes Phys. (Springer) **562** (2001) 195–212, gr-qc/0103067.
- [29] F.W. Hehl, A. Macías, *Metric-affine gauge theory of gravity II. Exact solutions*, Int. J. Mod. Phys. **D8** (1999) 399–416, gr-qc/9902076.
- [30] F.W. Hehl, J.D. McCrea, E.W. Mielke and Y. Ne’eman, *Metric-affine gauge theory of gravity: Field equations, Noether identities, world spinors, and breaking of dilation invariance*, Phys. Rept. **258** (1995) 1–171.
- [31] F. W. Hehl and Y. N. Obukhov, *Is a ‘hadronic’ shear current one of the sources in metric-affine gravity?*, 8th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Gravitation and Relativistic Field Theories (MG 8), Jerusalem, Israel, 22-27 June 1997, gr-qc/9712089.
- [32] F.W. Hehl and Yu.N. Obukhov, *Foundations of Classical Electrodynamics: Charge, Flux, and Metric* (Birkhäuser: Boston, MA, 2003).
- [33] C. Heinicke and F.W. Hehl, *Computer algebra in gravity*, 5 pages in: J. Grabmeier, E. Kaltofen, and V. Weispfenning, eds. *Computer Algebra Handbook: Foundations, Applications, Systems* (Springer: Berlin, 2003).
- [34] J.K. Ho, D.-C. Chern, and J.M. Nester, *Some spherically symmetric exact solutions of the metric-affine gravity theory*, Chinese J. Physics (Taipei) **35** (1997) 640–650.
- [35] Y. Itin, *Coframe energy-momentum current. Algebraic properties*, Gen. Rel. Grav. **34** (2002)

- 1819–1837, [gr-qc/0111087](#).
- [36] T. Jacobson, D. Mattingly, *Gravity with a dynamical preferred frame*, Phys. Rev. **D64** (2001) 024028 (9 pages), [gr-qc/0007031](#).
 - [37] T. Jacobson and D. Mattingly, *Einstein-aether waves*, Phys. Rev. **D70** (2004) 024003 (5 pages), [gr-qc/0402005](#). In this paper J&M finally set up the theory and give it a name, see also the references given there. We consider this as their central paper on the Einstein-aether.
 - [38] J. Julve, A. Lopez-Pinto, A. Tiemblo and R. Tresguerres, *Nonlinear gauge foundation of affine theories of gravity with fermionic matter*, Aug 1995. In *Sardanashvily, G.A. (ed.): New frontiers in gravitation* pp. 115–144.
 - [39] A.D. King and D. Vassiliev, *Torsion waves in metric-affine field theory*, Class. Quantum Grav. **18** (2001) 2317–2329.
 - [40] I. Kirsch, *A Higgs mechanism for gravity*, 14 pages, [hep-th/0503024](#) (2005).
 - [41] V.A. Kostelecky, *Gravity, Lorentz violation, and the Standard Model*, Phys. Rev. **D69** (2004) 105009 (20 pages), [hep-th/0312310](#).
 - [42] V.A. Kostelecky and S. Samuel, *Gravitational phenomenology in higher dimensional theories and strings*, Phys. Rev. **D40** (1989) 1886–1903.
 - [43] R. Kuhfuss and J. Nitsch, *Propagating modes in gauge field theories of gravity*, Gen. Relat. Grav. **18** (1986) 1207–1227.
 - [44] C. Lämmerzahl, *Constraints on space-time torsion from Hughes-Drever experiments*, Phys. Lett. **A228** (1997) 223–231, [gr-qc/9704047](#).
 - [45] C. Lämmerzahl, *Special relativity and Lorentz invariance*, Ann. Physik (Leipzig) **14** (2005) 71–102.
 - [46] E.A. Lim, *Can we see Lorentz-violating vector fields in the CMB?*, Phys. Rev. **D71** (2005) 063504 (16 pages), [astro-ph/0407437](#).
 - [47] A. Lopez-Pinto, A. Tiemblo and R. Tresguerres, *Non-Linear Affine Embedding of the Dirac Field from the Multiplicity-Free $SL(4, R)$ Unirreps*, Class. Quant. Grav. **13** (1996) 2255–2266.
 - [48] E.A. Lord, *A Unified approach to the gauging of space-time and internal symmetries*, Gen. Rel. Grav. **19** (1987) 983–1002.
 - [49] E.A. Lord and P. Goswami, *Gauge theory of a group of diffeomorphisms. I. General principles*, J. Math. Phys. **27** (1986) 2415–2422; ... *II. The conformal and de Sitter groups*, (Lord alone) J. Math. Phys. **27** (1986) 3051–3054; ... *III. The fiber bundle description*, J. Math. Phys. **29**

- (1988) 258–267.
- [50] A. Macías, C. Lämmerzahl and A. García, *A class of colliding waves in metric-affine gravity, nonmetricity and torsion shock waves*, J. Math. Phys. **41** (2000) 6369–6380, [gr-qc/0006007](#).
 - [51] D. Mattingly, *Modern tests of Lorentz invariance*, 65 pages, [gr-qc/0502097](#) (2005).
 - [52] E.W. Mielke: *Geometrodynamics of Gauge Fields* — On the geometry of Yang-Mills and gravitational gauge theories (Akademie-Verlag, Berlin 1987).
 - [53] E.W. Mielke, *Affine generalization of the Komar complex of general relativity*, Phys. Rev. **D63** (2001) 044018 (4 pages).
 - [54] A.V. Minkevich and Yu.G. Vasilevski, *Spherically-symmetric gravitational fields in the metric-affine gauge theory of gravity*, 17 pages, [gr-qc/0301098](#) (2003).
 - [55] H. Müller, S. Herrmann, A. Saenz, A. Peters and C. Lämmerzahl, *Tests of Lorentz invariance using hydrogen molecules*, Phys. Rev. **D70** (2004) 076004 (9 pages).
 - [56] Y. Ne’eman, F.W. Hehl, *Test matter in a spacetime with nonmetricity*, Class. Quant. Grav. **14** (1997) A251–A260, [gr-qc/9604047](#).
 - [57] J.M. Nester, *Gravity, torsion and gauge theory*, in: *Introduction to Kaluza–Klein theories*, H.C. Lee, ed. (World Scientific, Singapore 1984) pp. 83–115.
 - [58] Yu.N. Obukhov, *Plane waves in metric-affine gravity*, talk at a meeting of the German Physical Society (DPG) in Ulm, Germany, March 2004. To be published.
 - [59] Yu.N. Obukhov, E.J. Vlachynsky, W. Esser, and F.W. Hehl, *Effective Einstein theory from metric-affine gravity models via irreducible decompositions*, Phys. Rev. **D56** (1997) 7769–7778, [gr-qc/9705039](#).
 - [60] Yu.N. Obukhov, E.J. Vlachynsky, W. Esser, R. Tresguerres, and F.W. Hehl, *An exact solution of the metric-affine gauge theory with dilation, shear, and spin charges*, Phys. Lett. **A220** (1996) 1–9, [gr-qc/9604027](#).
 - [61] L. O’Raifeartaigh, *The Dawning of Gauge Theory* (Princeton University Press: Princeton, NJ, 1997).
 - [62] R. Percacci, *The Higgs phenomenon in quantum gravity*, Nucl. Phys. **B353** (1991) 271–290.
 - [63] O. Preuss, *Astronomical tests of the Einstein equivalence principle*. Ph.D. Thesis, University of Bielefeld (November 2002) 125 pages, [gr-qc/0305083](#).
 - [64] O. Preuss, M.P. Haugan, S.K. Solanki, S. Jordan, *Astronomical search for evidence of new physics: Limits of gravity-induced birefringence from the magnetic white dwarf RE J0317-853*,

- Phys. Rev. **D70** (2004) 067101 (4 pages), [gr-qc/0405068](#).
- [65] D. Puetzfeld, *A cosmological model in Weyl-Cartan spacetime: I. Field equations and solutions, II. Magnitude-redshift relation*, Class. Quant. Grav. **19** (2002) 3263–3280, 4463–4482, [gr-qc/0111014](#), [gr-qc/0205052](#).
 - [66] D. Puetzfeld, *Status on non-Riemannian cosmology*, [gr-qc/0404119](#), 10 pages (2004).
 - [67] D. Puetzfeld, X. Chen, *Testing non-standard cosmological models with supernovae*, [gr-qc/0402026](#) (2004).
 - [68] D. Puetzfeld and R. Tresguerres, *A cosmological model in Weyl-Cartan spacetime*, Class. Quant. Grav. **18** (2001) 677–694, [gr-qc/0101050](#).
 - [69] E. Schrödinger, *Space-Time Structure* (Cambridge University Press: Cambridge, 1954).
 - [70] E. Sezgin and P. van Nieuwenhuizen, *New ghost-free Lagrangians with propagating torsion*, Phys. Rev. **D21** (1980) 3269–3280.
 - [71] J. Socorro, A. Macias and F. W. Hehl, *Computer algebra in gravity: Reduce-Excalc programs for (non-)Riemannian spacetimes. I*, Comput. Phys. Comm. **115** (1998) 264–283, [gr-qc/9804068](#).
 - [72] S.K. Solanki, O. Preuss, M.P. Haugan, et al., *Solar constraints on new couplings between electromagnetism and gravity*, Phys. Rev. **D69** (2004) 062001 (11 pages), [gr-qc/0402055](#).
 - [73] W. Thirring, *Classical Mathematical Physics – Dynamical Systems and Field Theories*, 3rd ed. (Springer: New York, 1997).
 - [74] M.E. Tobar, P. Wolf, A. Fowler and J.G. Hartnett, *New methods of testing Lorentz violation in electrodynamics*, Phys. Rev. **D71** (2005) 025004 (15 pages), [hep-ph/0408006](#).
 - [75] A. Trautman, *The Einstein-Cartan theory*, in: *Encyclopedia of Mathematical Physics*, J.P. Francoise et al. (eds.). Elsevier, Oxford, 13 pages, to be published (2005) [<http://www.fuw.edu.pl/~amt/ect.pdf>].
 - [76] R. Tresguerres, *Exact vacuum solutions of 4-dimensional metric-affine gauge theories of gravitation*, Z. Phys. **C65** (1995) 347–354.
 - [77] R. Tresguerres, *Exact static vacuum solution of four-dimensional metric-affine gravity with nontrivial torsion*, Phys. Lett. **A200** (1995) 405–410.
 - [78] R. Tresguerres, *Unified description of interactions in terms of composite fiber bundles*, Phys. Rev. **D66** (2002) 064025 (15 pages).
 - [79] R. Tresguerres and E. W. Mielke, *Gravitational Goldstone fields from affine gauge theory*,

- Phys. Rev. **D62** (2000) 044004 (7 pages), [gr-qc/0007072](#).
- [80] R.W. Tucker and C. Wang, *Black holes with Weyl charge and non-Riemannian waves*, Class. Quantum Grav. **12** (1995) 2587–2605.
 - [81] R.W. Tucker and C. Wang, *Dark matter gravitational interactions*, Class. Quantum Grav. **15** (1998) 933–954, [gr-qc/9612019](#).
 - [82] D. Vassiliev, *Pseudoinstantons in metric-affine field theory*, Gen. Relat. Grav. **34** (2002) 1239–1265.
 - [83] D. Vassiliev, *Quadratic metric-affine gravity*, Ann. Physik (Leipzig) **14** (2005) 231–252, [gr-qc/0304028](#).
 - [84] H. Weyl, *Gravitation und Elektrizität*, Sitzungsber. Königl. Preuss. Akad. Wiss. (Berlin), pp.265–480 (1918).
 - [85] Y.-H. Wu, *Quasilocal energy-momentum in Metric Affine Gravity*, Master thesis. National Central University, Chung-Li, Taiwan, ROC (June 2001) 66 pages.
 - [86] Y.-H. Wu, J.M. Nester, and C.-M. Chen, *Quasilocal energy-momentum in metric affine gravity: Asymptotic values for spherically symmetric solutions*, preprint National Central University, Chung-li, Taiwan, ROC, to be published (2003/04).
 - [87] Y.-H. Wu, C.-M. Chen, and J.M. Nester, *Quasilocal energy-momentum in metric affine gravity I: Dilaton type solution*, preprint National Central University, Chung-li, Taiwan, ROC, to be published (2003).

=====